



**Michigan
Technological
University**

Michigan Technological University
Digital Commons @ Michigan Tech

Department of Chemistry Publications

Department of Chemistry

11-4-2011

Controlled Knoevenagel reactions of methyl groups of 1,3,5,7-tetramethyl BODIPY dyes for unique BODIPY dyes

Shilei Zhu

Michigan Technological University

Jingtuo Zhang

Michigan Technological University

Giri Vegesna

Michigan Technological University

Ashutosh Tiwari

Michigan Technological University

Fen-Tair Luo

Academia Sinica

See next page for additional authors


Follow this and additional works at: <https://digitalcommons.mtu.edu/chemistry-fp>

 Part of the [Chemistry Commons](#)

Recommended Citation

Zhu, S., Zhang, J., Vegesna, G., Tiwari, A., Luo, F., Zeller, M., Luck, R., Li, H., Green, S., & Liu, H. (2011). Controlled Knoevenagel reactions of methyl groups of 1,3,5,7-tetramethyl BODIPY dyes for unique BODIPY dyes. *RSC Advances*, 2, 404-407. <http://dx.doi.org/10.1039/C1RA00678A>
Retrieved from: <https://digitalcommons.mtu.edu/chemistry-fp/113>

Follow this and additional works at: <https://digitalcommons.mtu.edu/chemistry-fp>

 Part of the [Chemistry Commons](#)

Authors

Shilei Zhu, Jingtuo Zhang, Giri Vegesna, Ashutosh Tiwari, Fen-Tair Luo, Matthias Zeller, Rudy Luck, Haihua Li, Sarah Green, and Haiying Liu

Cite this: *RSC Advances*, 2012, 2, 404–407

www.rsc.org/advances

COMMUNICATION

Controlled Knoevenagel reactions of methyl groups of 1,3,5,7-tetramethyl BODIPY dyes for unique BODIPY dyes†

Shilei Zhu,^a Jingtuo Zhang,^a Giri Vegesna,^a Ashutosh Tiwari,^a Fen-Tair Luo,^b Matthias Zeller,^c Rudy Luck,^a Haihua Li,^d Sarah Green^a and Haiying Liu*^a

Received 5th September 2011, Accepted 10th October 2011

DOI: 10.1039/c1ra00678a

Formyl groups at 6- and 2,6-positions initiated Knoevenagel reactions of the methyl groups at the 7, and 1,7-positions of 1,3,5,7-tetramethyl BODIPY dyes with aromatic aldehydes. Formation of vinyl bonds at the 7-, and 1,7-positions facilitates further Knoevenagel reactions of the methyl groups at the 3,5-positions. This approach offers fast, facile and versatile ways to prepare potential novel building blocks of BODIPY dyes for conjugated oligomers, dendrimers, and highly water-soluble, near-infrared emissive sensing materials.

BODIPY (4,4-difluoro-4-bora-3a,4a-diaza-s-indacene) dyes have received considerable attention recently because they possess many distinctive and desirable properties such as high extinction coefficients, narrow absorption and emission bands, high quantum efficiencies of fluorescence, relative insensitivity to environmental perturbations, and high resistance to photobleaching.¹ BODIPY dyes typically absorb light in the visible region and emit fluorescence between 470 and 550 nm. Tuning of their optical properties for absorption and emission maxima in the far-red and near-infrared regions has been achieved by functionalizing BODIPY cores at the *meso*-, 2,6- and 3,5-positions, fusing some aromatic rings to the BODIPY core, and replacing pyrrole by isoindole.^{1–3} Among these approaches, monostyryl- and distyryl modifications *via* the Knoevenagel reaction are facile and versatile routes to effectively tune BODIPY dyes for near-infrared emission. Methyl groups at the 3,5-positions of BODIPY dyes are active in the Knoevenagel reaction, often resulting in 3-monostyryl- and 3,5-distyryl-BODIPY dyes.^{1–6} Recently, the Akkaya group prepared 3,5,7-tristyryl- and 1,3,5,7-tetrastryryl-BODIPY dyes by activating methyl groups at the

1,7-positions *via* the introduction of phenylethynyl groups at the 2,6-positions.⁷ Ziessel's group successfully prepared 1,3,5,7-tetrastryryl-BODIPY dyes by using a high temperature of 140 °C.⁸ However, methyl groups at the 3,5-positions in these reported approaches are more reactive and undergo Knoevenagel reactions faster than those at the 1,7-positions.^{7,8}

In this communication, we report a simple and efficient approach to control the Knoevenagel reaction sequence of methyl groups at the 1,3,5,7-positions of BODIPY dyes. The method is utilized to prepare novel building blocks of monostyryl, distyryl-, tristyryl- and tetrastryryl-BODIPY dyes with unique chemical structures. To achieve this goal we employed electron withdrawing formyl groups at the 6-, and 2,6-positions of BODIPY dyes to initiate Knoevenagel reactions of methyl groups at the 7-, and the 1,7-positions with aromatic aldehyde derivatives. These formyl groups may provide steric hindrance and prevent initial Knoevenagel reaction of methyl groups at the 5, and the 3,5-positions with aromatic aldehydes, resulting in 2-iodo-6-formyl-3,7-distyryl-BODIPY dye, 2,6-diformyl-1-monostyryl-BODIPY dye and, 2,6-diformyl-1,7-distyryl-BODIPY dye. Formation of vinyl bonds at the 7-, and 1,7-positions facilitates further Knoevenagel reaction of methyl groups at the 3, and the 3,5-positions with the aromatic aldehydes, producing 2-iodo-6-formyl-3,5,7-tristyryl-BODIPY dye, 2,6-diformyl-1,3,7-tristyryl-BODIPY dye, and 2,6-diformyl-1,3,5,7-tetrastryryl-BODIPY dye. Branched oligo(ethylene glycol)methyl ether residues were used to facilitate water solubility of 1,7-distyryl and 1,3,5,7-tetrastryryl-BODIPY dyes. These BODIPY dyes can serve as novel potential building blocks for BODIPY-based conjugated oligomers, dendrimers, and highly water-soluble near-infrared emissive sensing materials.

We chose reported BODIPY dye (**2**) to illustrate the feasibility of our unique approach. A formyl group was introduced to BODIPY dye **2** at the 6-position *via* the Vilsmeier–Haack reaction, affording 6-formyl-BODIPY dye (**3**). Iodination of BODIPY dye **3** at the 2-position yields 2-iodo-6-formyl-BODIPY dye (**4**). A three-hour Knoevenagel condensation of BODIPY dye **4** with excess 3,4,5-trimethoxybenzaldehyde (**5**) afforded 2-iodo-6-formyl-3-monostyryl-BODIPY dye (**A**), 2-iodo-6-formyl-3,7-distyryl-BODIPY dye (**B**) and 2-iodo-6-formyl-3,5,7-tristyryl-BODIPY dye (**C**) in diluted solution (Scheme 1). Introduction of an iodo group at the 2-position helps activate methyl groups at the 3,7-positions of BODIPY dye **4**. These heterobifunctional BODIPY dyes are expected to be potential and useful building blocks for BODIPY-based donor– π -acceptor

^aDepartment of Chemistry, Michigan Technological University, 1400 Townsend Drive, Houghton, MI, 49931, USA. E-mail: hylu@mtu.edu; Fax: +1 9064872061; Tel: +1 9064873451

^bInstitute of Chemistry, Academia Sinica, Taipei, Taiwan, 11529, Republic of China

^cDepartment of Chemistry, Youngstown State University, Youngstown, OH, 44555, USA

^dUniversity Libraries, University of Alabama, Tuscaloosa, AL, 35487, USA

† Electronic supplementary information (ESI) available: Synthesis, characterization and optical properties of BODIPY dyes, and crystal structures of BODIPY dyes **4** and **6**. CCDC reference numbers 826761 and 826762. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c1ra00678a

Table 1 Absorption and emission peaks, fluorescence quantum yields (Φ_f), and absorption coefficients (ϵ_{\max}) of BODIPY dyes (2–4, 6, A–D_a, E_a, F_a and G_a) in methylene chloride solution and BODIPY dyes (E_b, G_b) aqueous solution. Fluorescence quantum yields of BODIPY dyes were calculated by using fluorescein excited at 490 nm in 0.1 N NaOH as the reference quantum efficiency ($\Phi_f = 85\%$)^{13–16}

BODIPY dye	$\lambda_{\text{abs}}/\text{nm}$	$\lambda_{\text{em}}/\text{nm}$	Φ_f (%)	$\epsilon_{\max}/10^4 \text{ M}^{-1}\text{cm}^{-1}$
2	500	510	80	8.86
3	497	508	65.5	8.24
4	518	530	1.5	8.26
6	504	514	48.7	12.67
A	582	615	3.5	7.99
B	617	660	2.1	3.97
C	665	698	0.9	5.86
D _a	537	621	0.5	2.53
E _a	576	673	0.2	3.69
E _b	557	695	0.1	1.29
F _a	634	686	0.1	6.06
G _a	688	722	0.4	7.27
G _b	661	710	0.1	4.58

affect π -conjugation of BODIPY dyes. 2,6-Diformyl-1,3,5,7-tetraaryl-BODIPY dye (G_b) is the first reported water-soluble neutral near-infrared emissive BODIPY dye to the best of our knowledge. All new BODIPY dyes (A–G) show low fluorescence quantum yields. Fluorescence of BODIPY dyes A–C is expected to be quenched both by the heavy element iodine and an aldehyde group, while BODIPY dyes D–G are expected to be quenched by two aldehyde groups.

It seems that the formyl groups quench the fluorescence of BODIPY dyes with extended π -conjugation systems more efficiently than that of typical BODIPY dyes (Table 1). In order to test our hypothesis that formyl groups quench the fluorescence of BODIPY dyes, we measured the optical properties of BODIPY dyes (E_b and G_b) in the presence of cysteine, that can react with aldehyde groups resulting in thiazolidine formation.¹² The presence of a 500-fold equivalent of cysteine in ten hours significantly enhances fluorescence intensity (up to 210 times) of the BODIPY dye E_b, and causes blue shifts in both absorption and emission by 64 nm and 96 nm (Fig. 3 and 4), respectively as the reaction of aldehyde groups with cysteine resulted in thiazolidine formation¹² and reduced π -conjugation of the BODIPY dyes. The presence of cysteine also causes color changes of the aqueous solutions of BODIPY dye E_b from blue to purple, and colorless to red in the absence (Fig. 5a) and presence of a transilluminator, respectively (see Fig. 44 in the ESI†). The presence of 500-fold equivalents of cysteine resulted in a 60-fold increase in




Fig. 1 Normalized absorption spectra of BODIPY dyes (4, 6, A–C, D_a–G_a) in methylene chloride.




Fig. 2 Normalized fluorescence spectra of BODIPY dyes (4, 6, A–C, D_a–G_a) in methylene chloride.




Fig. 3 Normalized absorption spectra of BODIPY dye E_b (20 μM) in 0.01 M PBS buffer pH 7.4 in the absence and presence of L-cysteine (10 mM) at different times.

fluorescence intensity of BODIPY dye G_b, causing the disappearance of an absorption peak at 400 nm and leading to blue shifts in both absorption and emission by 37 nm and 33 nm, respectively (see Fig. 41 and 42 in the ESI†). The presence of cysteine changes the color of the aqueous solution of BODIPY dye G_b from green to blue (Fig. 5b), and colorless to red in the absence and presence of a transilluminator, respectively (see Fig. S45 in the ESI†). These results demonstrate that the presence of the formyl groups quenches the dye fluorescence.




Fig. 4 Normalized fluorescence spectra of BODIPY dye E_b (20 μM) in 0.01 M PBS buffer pH 7.4 in the absence and presence of L-cysteine (10 mM) at different times. Excitation wavelength was at 470 nm.




Fig. 5 (a) Visualized images of BODIPY dye E_b ($20\ \mu\text{M}$) in 0.01M PBS buffer pH 7.4 in the absence and presence of L-cysteine (10 mM). (b) Visualized images of BODIPY dye G_b ($20\ \mu\text{M}$) in 0.01M PBS buffer pH 7.4 in the absence and presence of L-cysteine (10 mM).




Fig. 6 Fluorescence response of BODIPY dye G_b ($10\ \mu\text{M}$) to various amino acids (5 mM) at 677 nm. Conditions: 0.01M PBS buffer pH 7.4 solution. $\lambda_{\text{ex}} = 620\ \text{nm}$. Black bars represent the addition of various amino acids to G_b solution. White bars represent the further addition of cysteine (5 mM) to the above solutions, respectively.

The selective response of BODIPY dye G_b to cysteine was tested against the presence of other amino acids. The presence of 500 equivalents of other amino acids does not cause any significant changes in fluorescence intensity and absorption of BODIPY dye G_b (Fig. 6 and Fig. S48 in the ESI†). These BODIPY dyes may offer the effective colorimetric and fluorimetric detection of cysteine and

might have potential application for the sensing and labeling of N-terminal cysteine residues in peptides or proteins.

In conclusion, we have developed an effective approach to control the Knoevenagel reaction sequence of methyl groups of 1,3,5,7-tetramethyl BODIPY dyes by placing formyl groups at the 2- and the 2,6-positions. This allows for the preparation of novel near-infrared emissive BODIPY dyes, which can serve as potential useful building blocks for oligomers, dendrimers and sensing and imaging materials.

Acknowledgements

H.Y. Liu acknowledges an NIH SBIR grant (contract number: 1R43CA134039-01) and NSF grant (award number: 1048655) for financial support and A. Tiwari acknowledges the ALS Therapy Alliance for financial support. The X-ray diffractometer was funded by NSF Grant 0087210, Ohio Board of Regents Grant CAP-491, and by Youngstown State University.

References

- 1 A. Loudet and K. Burgess, *Chem. Rev.*, 2007, **107**, 4891–4932.
- 2 R. Ziessel, G. Ulrich and A. Harriman, *New J. Chem.*, 2007, **31**, 496–501.
- 3 G. Ulrich, R. Ziessel and A. Harriman, *Angew. Chem., Int. Ed.*, 2008, **47**, 1184–1201.
- 4 S. Zhu, J. Zhang, G. K. Vegesna, R. Pandey, F. T. Luo, S. A. Green and H. Liu, *Chem. Commun.*, 2011, **47**, 3508–3510.
- 5 S. Zhu, J. Zhang, G. Vegesna, F. T. Luo, S. A. Green and H. Liu, *Org. Lett.*, 2011, **13**, 438–441.
- 6 Z. Ekmekci, M. D. Yilmaz and E. U. Akkaya, *Org. Lett.*, 2008, **10**, 461–464.
- 7 O. Buyukcakir, O. A. Bozdemir, S. Kolemen, S. Erbas and E. U. Akkaya, *Org. Lett.*, 2009, **11**, 4644–4647.
- 8 T. Bura, P. Retailleau, G. Ulrich and R. Ziessel, *J. Org. Chem.*, 2011, **76**, 1109–1117.
- 9 L. Jiao, C. Yu, J. Li, Z. Wang, M. Wu and E. Hao, *J. Org. Chem.*, 2009, **74**, 7525–7528.
- 10 O. A. Bozdemir, R. Guliyev, O. Buyukcakir, S. Selcuk, S. Kolemen, G. Gulseren, T. Nalbantoglu, H. Boyaci and E. U. Akkaya, *J. Am. Chem. Soc.*, 2010, **132**, 8029–8036.
- 11 D. Buccella, J. A. Horowitz and S. J. Lippard, *J. Am. Chem. Soc.*, 2011, **133**, 4101–4114.
- 12 O. Rusin, N. N. St Luce, R. A. Agbaria, J. O. Escobedo, S. Jiang, I. M. Warner, F. B. Dawan, K. Lian and R. M. Strongin, *J. Am. Chem. Soc.*, 2004, **126**, 438–439.
- 13 C. A. Parker and W. T. Rees, *Analyst*, 1960, **85**, 587–600.
- 14 T. Yogo, Y. Urano, Y. Ishitsuka, F. Maniwa and T. Nagano, *J. Am. Chem. Soc.*, 2005, **127**, 12162–12163.
- 15 S. Maruyama, K. Kikuchi, T. Hirano, Y. Urano and T. Nagano, *J. Am. Chem. Soc.*, 2002, **124**, 10650–10651.
- 16 Y. Gabe, Y. Urano, K. Kikuchi, H. Kojima and T. Nagano, *J. Am. Chem. Soc.*, 2004, **126**, 3357–3367.