



**Michigan
Technological
University**

Michigan Technological University
Digital Commons @ Michigan Tech

Michigan Tech Publications

3-24-2022

Do 2H and 18O in leaf water reflect environmental drivers differently?

Lucas A. Cernusak
James Cook University

Adrià Barbeta
Universitat de Barcelona

Rosemary T. Bush
Northwestern University

Rebekka Eichstaedt Bögelein
University of Trier

Juan Pedro Ferrio
Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA)

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.mtu.edu/michigantech-p>



Part of the [Forest Sciences Commons](#)

Recommended Citation

Cernusak, L. A., Barbeta, A., Bush, R. T., Eichstaedt Bögelein, R., Ferrio, J. P., Flanagan, L. B., Gessler, A., Martín-Gómez, P., Hirl, R. T., Kahmen, A., Keitel, C., Lai, C., Munksgaard, N. C., Nelson, D. B., Ogée, J., Roden, J. S., Schnyder, H., Voelker, S., Wang, L., Stuart-Williams, H., Wingate, L., Yu, W., Zhao, L., & Cuntz, M. (2022). Do 2H and 18O in leaf water reflect environmental drivers differently?. *The New phytologist*. <http://doi.org/10.1111/nph.18113>
Retrieved from: <https://digitalcommons.mtu.edu/michigantech-p/15906>

Follow this and additional works at: <https://digitalcommons.mtu.edu/michigantech-p>



Part of the [Forest Sciences Commons](#)

Authors























Lucas A. Cernusak, Adrià Barbeta, Rosemary T. Bush, Rebekka Eichstaedt Bögelein, Juan Pedro Ferrio, Lawrence B. Flanagan, Arthur Gessler, Paula Martín-Gómez, Regina T. Hirl, Ansgar Kahmen, Claudia Keitel, Chun-Ta Lai, Niels C. Munksgaard, Daniel B. Nelson, Jérôme Ogée, John S. Roden, Hans Schnyder, Steve L. Voelker, Lixin Wang, Hilary Stuart-Williams, Lisa Wingate, Wusheng Yu, Liangju Zhao, and Matthias Cuntz

Research review

Do ^2H and ^{18}O in leaf water reflect environmental drivers differently?

Author for correspondence:
Lucas A. Cernusak
Email: lucas.cernusak@jcu.edu.au

Received: 18 November 2021
Accepted: 6 February 2022

Lucas A. Cernusak¹ , Adrià Barbeta² , Rosemary T. Bush³,
Rebekka Eichstaedt (Bögelein)⁴ , Juan Pedro Ferrio⁵ ,
Lawrence B. Flanagan⁶ , Arthur Gessler⁷ , Paula Martín-Gómez^{8,9} ,
Regina T. Hirl¹⁰, Ansgar Kahmen¹¹ , Claudia Keitel¹² , Chun-Ta Lai¹³ ,
Niels C. Munksgaard¹ , Daniel B. Nelson¹¹ , Jérôme Ogée⁸ ,
John S. Roden¹⁴ , Hans Schnyder¹⁰ , Steven L. Voelker¹⁵ ,
Lixin Wang¹⁶ , Hilary Stuart-Williams¹⁷ , Lisa Wingate⁸ ,
Wusheng Yu¹⁸ , Liangju Zhao¹⁹  and Matthias Cuntz²⁰ 

¹College of Science and Engineering, James Cook University, Cairns, Qld 4878, Australia; ²BEECA, Department of Evolutionary Biology, Ecology and Environmental Sciences, Universitat de Barcelona, Barcelona, Catalonia 08028, Spain; ³Department of Earth and Planetary Sciences, Northwestern University, Evanston, IL 60208, USA; ⁴Faculty of Regional and Environmental Sciences – Geobotany, University of Trier, Trier 54296, Germany; ⁵ARAID-Departamento de Sistemas Agrícolas, Forestales y Medio Ambiente, Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA), Zaragoza 50059, Spain; ⁶Department of Biological Sciences, University of Lethbridge, Lethbridge, AB T1K 3M4, Canada; ⁷WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Birmensdorf 8903, Switzerland; ⁸INRAE, Bordeaux Sciences Agro, UMR ISPA, Villenave d'Ornon 33140, France; ⁹Centre Tecnològic Forestal de Catalunya (CTFC), Solsona, Catalonia 25280, Spain; ¹⁰Technische Universität München, Lehrstuhl für Grünlandlehre, Freising-Weihenstephan 85354, Germany; ¹¹Department of Environmental Sciences–Botany, University of Basel, Basel 4056, Switzerland; ¹²School of Life and Environmental Sciences, Sydney Institute of Agriculture, The University of Sydney, Camden, NSW 2006, Australia; ¹³Department of Biology, San Diego State University, San Diego, CA 92182, USA; ¹⁴Department of Biology, Southern Oregon University, Ashland, OR 97520, USA; ¹⁵College of Forest Resources and Environmental Science, Michigan Technological University, Houghton, MI 49931, USA; ¹⁶Department of Earth Sciences, Indiana University–Purdue University Indianapolis, Indianapolis, IN 46202, USA; ¹⁷Research School of Biology, The Australian National University, Canberra, ACT 2601, Australia; ¹⁸Key Laboratory of Tibetan Environmental Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China; ¹⁹Shaanxi Key Laboratory of Earth Surface System and Environmental Carrying Capacity, College of Urban and Environmental Sciences, Northwest University, Xi'an 710127, China; ²⁰Université de Lorraine, AgroParisTech, INRAE, UMR Silva, Nancy 54000, France

Summary

We compiled hydrogen and oxygen stable isotope compositions ($\delta^2\text{H}$ and $\delta^{18}\text{O}$) of leaf water from multiple biomes to examine variations with environmental drivers. Leaf water $\delta^2\text{H}$ was more closely correlated with $\delta^2\text{H}$ of xylem water or atmospheric vapour, whereas leaf water $\delta^{18}\text{O}$ was more closely correlated with air relative humidity. This resulted from the larger proportional range for $\delta^2\text{H}$ of meteoric waters relative to the extent of leaf water evaporative enrichment compared with $\delta^{18}\text{O}$. We next expressed leaf water as isotopic enrichment above xylem water ($\Delta^2\text{H}$ and $\Delta^{18}\text{O}$) to remove the impact of xylem water isotopic variation. For $\Delta^2\text{H}$, leaf water still correlated with atmospheric vapour, whereas $\Delta^{18}\text{O}$ showed no such correlation. This was explained by covariance between air relative humidity and the $\Delta^{18}\text{O}$ of atmospheric vapour. This is consistent with a previously observed diurnal correlation between air relative humidity and the deuterium excess of atmospheric vapour across a range of ecosystems. We

New Phytologist (2022)
doi: 10.1111/nph.18113

Key words: evaporative enrichment, isotopic biomarker, leaf water, relative humidity, stable isotopes.

conclude that ^2H and ^{18}O in leaf water do indeed reflect the balance of environmental drivers differently; our results have implications for understanding isotopic effects associated with water cycling in terrestrial ecosystems and for inferring environmental change from isotopic biomarkers that act as proxies for leaf water.

Introduction

The stable isotope composition of hydrogen and oxygen in leaf water varies throughout the day, among plants within a site and across environmental gradients (Zundel *et al.*, 1978; Flanagan *et al.*, 1991a; Cernusak *et al.*, 2002, 2016; Lai *et al.*, 2008; West *et al.*, 2008). Leaf water becomes enriched in the heavy isotopes ^2H and ^{18}O compared with the water entering the roots as a result of evaporative isotopic fractionation during transpiration (Gonfiantini *et al.*, 1965). There is also isotopic exchange between water vapour in the atmosphere and that in the leaf (Craig & Gordon, 1965); notably, this continues even if transpiration has ceased under a saturated atmosphere (Welp *et al.*, 2008; Kim & Lee, 2011; Helliker, 2014; Goldsmith *et al.*, 2017). Furthermore, the distribution of isotope enrichment within the leaf can vary as a function of leaf anatomy and physiology (Yakir *et al.*, 1989; Gan *et al.*, 2002; Holloway-Phillips *et al.*, 2016; Barbour *et al.*, 2021). Therefore, the stable isotope composition of leaf water provides an information-rich isotopic signal that can be applied across a broad range of disciplines (Yakir, 1998). Interest in understanding leaf water stable isotope composition has been further motivated by recognition that leaf water is the starting point for isotope signals in plant organic compounds such as sucrose, starch, cellulose, lignin, leaf waxes (Yakir, 1992; Farquhar *et al.*, 1998; Barbour, 2007; Lehmann *et al.*, 2020). Leaf water isotopic signals can even be reflected in the bones and teeth of herbivores, such as kangaroos (Ayliffe & Chivas, 1990; Faith, 2018).

Models of leaf water stable isotope composition have been developed over several decades and typically perform reasonably well at explaining observed leaf water isotopic variation (Dongmann *et al.*, 1974; Flanagan *et al.*, 1991b; Roden & Ehleringer, 1999; Farquhar & Cernusak, 2005; Cuntz *et al.*, 2007; Ogée *et al.*, 2007). However, some questions about subtler aspects of leaf water isotopic composition remain (Cernusak *et al.*, 2016). One such question is whether stable isotopes of hydrogen and oxygen reflect differently the balance of environmental and physiological drivers that lead to variation in leaf water stable isotopes.

Models of leaf water isotopic composition do not differentiate between hydrogen and oxygen in their general formulation; the major mechanisms that cause leaf water to change isotopically are common to both elements. However, the magnitudes of the fractionation factors associated with the mechanisms do differ. This is also true for meteoric waters, such that the relative extent of variation in the isotopic composition of plant source water and atmospheric vapour across the landscape is different between ^2H and ^{18}O ; on average, there is a *c.* 8‰ change in $\delta^2\text{H}$ for a given 1‰ change in $\delta^{18}\text{O}$ (Craig, 1961; Rozanski *et al.*, 1993). Movement of the two isotopologues H_2^{18}O and ^2HHO within the leaf may also vary, for example due to different diffusivities in water (Cuntz *et al.*, 2007), in air (Barbour *et al.*, 2017), and potentially across

membranes (Mamonov *et al.*, 2007) and there can be different extents of exchange with organic molecules (Yakir, 1992; Chen *et al.*, 2020). Here, we aimed to assess whether hydrogen and oxygen stable isotopes in leaf water respond differently to the environment, to better understand whether $\delta^2\text{H}$ and $\delta^{18}\text{O}$ in organic matter proxies capture environmental signals differently.

To do this, we compiled datasets that provided measurements under natural conditions of both $\delta^2\text{H}$ and $\delta^{18}\text{O}$ in leaf water, xylem water and atmospheric vapour, along with concurrent measurements of air temperature and relative humidity. Table 1 provides a summary of data sources. Within each dataset, we averaged individual observations, such that each row of data in the compiled dataset represents a mean value for a given species by site by time combination. In total, the dataset contained 546 such rows. The geographic range of the combined dataset covered more than 100° of latitude and more than 3000 m of elevation (Table 1). We limited the dataset to daytime observations, as it is primarily during photosynthesis that leaf water signals are incorporated into organic compounds. This also helped to avoid issues of nonsteady state leaf water enrichment at night (Cernusak *et al.*, 2002, 2005; Seibt *et al.*, 2006). We note that it has recently been shown that extraction of stem xylem water for isotopic analysis can be accompanied by an offset in $\delta^2\text{H}$ from the water that is likely to have been taken up by the roots (Zhao *et al.*, 2016; Chen *et al.*, 2020; Barbeta *et al.*, 2022). We did not attempt to apply a correction for this offset as we lacked a basis on which to make the correction that could be applied across the compiled dataset.

$\delta^2\text{H}$ and $\delta^{18}\text{O}$ of leaf water, xylem water and vapour

The Craig–Gordon equation (Craig & Gordon, 1965) forms the basic building block for models of leaf water isotopic composition and provides a convenient entry point for examining the environmental drivers of leaf water $\delta^2\text{H}$ and $\delta^{18}\text{O}$. The Craig–Gordon equation can be approximated as:

$$\delta_e \approx \delta_s + \epsilon^+ + \epsilon_k + (\delta_v - \delta_s - \epsilon_k)h \quad \text{Eqn 1}$$

(δ_e , predicted $\delta^2\text{H}$ or $\delta^{18}\text{O}$ at the evaporative sites within leaves; δ_s , $\delta^2\text{H}$ or $\delta^{18}\text{O}$ of source water, which we equated in our dataset to xylem water; ϵ^+ , equilibrium fractionation between liquid and vapour; ϵ_k , kinetic fractionation during diffusion through the stomata and boundary layer; δ_v , $\delta^2\text{H}$ or $\delta^{18}\text{O}$ of atmospheric vapour and h , w_a/w_i , the water vapour mole fraction in the air outside the leaf boundary layer divided by that at the evaporative sites inside the leaf substomatal cavity). The w_i is typically assumed to be saturated at leaf temperature, although recent evidence has suggested that it may be less than saturated at times (Cernusak *et al.*, 2018, 2019; Buckley & Sack, 2019; Holloway-Phillips *et al.*, 2019). If w_i is saturated and leaf temperature is equal to air

Table 1 Datasets and associated site information for the data compilation presented in this paper.

Dataset	Site	Latitude	Longitude	Elevation (m)	MAP (mm)	MAT (°C)	Vegetation type	References
Western_USA_Roden	Cascade_Heads	45.03	−123.91	14	2410	10.7	Forest	(Roden & Ehleringer, 2000a,b)
	Bill_Williams_River	34.26	−114.03	150	97	23.8	Woodland	
	Weber_River	41.13	−111.90	1450	510	10.6	Woodland	
	Red_Butte_Canyon	40.78	−111.80	1790	700	10.1	Woodland	
	Big_Cottonwood	40.62	−111.73	1987	840	9.4	Woodland	
Washington_USA_Lai	Wind_River	45.82	−121.95	371	2467	8.7	Forest	(Lai & Ehleringer, 2011)
Utah_USA_Flanagan	Coral_Pink	37.04	−112.72	1855	380	10.5	Woodland	(Flanagan <i>et al.</i> , 1993)
Tibetan_Plateau_Yu	Lhasa	29.65	91.03	3658	460	8.4	Grassland	W. Yu, unpublished
Qld_Aus_Munksgaard	Cairns	−16.79	145.69	30	2000	25.0	Forest/Woodland	(Munksgaard <i>et al.</i> , 2017)
	Tinaroo	−17.17	145.54	680	1400	22.0	Forest/Woodland	
	Herberton	−17.34	145.42	918	1150	19.0	Woodland	
	Wild_River	−17.65	145.28	860	950	21.0	Woodland	
	Mount_Garnet	−17.67	145.10	660	800	24.0	Woodland	
NW_China_Zhao	Pailugou_2900	38.54	100.30	2900	369.2	0.7	Forest	(Zhao <i>et al.</i> , 2014)
	Pailugou_2700	38.55	100.29	2780	369.2	0.7	Forest	
	Riparian	42.02	101.23	930	34.9	8.9	Woodland	
	Gobi	42.27	101.12	906	34.9	8.9	Woodland	
NT_Aus_Cernusak	Alice_Springs	−23.70	133.83	598	276	21.0	Woodland	(Kahmen <i>et al.</i> , 2013a; Cernusak <i>et al.</i> , 2016)
	Tennant_Creek	−19.65	134.16	365	454	25.9	Woodland	
	Elliot	−17.50	133.51	234	604	26.8	Woodland	
	Katherine	−14.48	132.36	143	1140	27.2	Woodland	
	Darwin	−12.44	130.88	33	1736	27.6	Woodland	
NSW_Aus_Twining	Tumbarumba	−35.66	148.15	1249	1900	9.6	Forest	(Twining <i>et al.</i> , 2006)
Hawaii_USA_Kahmen	MLM_1	19.69	155.20	683	5676	18.4	Forest	(Kahmen <i>et al.</i> , 2011)
	MLM_3	19.66	155.47	2061	2000	11.3	Forest/Woodland	
	MLM_4	19.59	155.45	2465	1500	9.9	Forest/Woodland	
	MLM_5	19.83	155.82	694	500	20.0	Forest/Woodland	
Greenland_Bush	Kangerlussuaq	67.02	−50.70	50	140	−5.7	Grassland	(Bush <i>et al.</i> , 2017)
Germany_Hirl	Grünschaige	48.40	11.75	448	743	9.3	Grassland	(Hirl <i>et al.</i> , 2019)
Germany_Bögelein	Palatinate	49.28	7.81	550	1067	7.9	Forest	(Bögelein <i>et al.</i> , 2017)
France_Wingate	LeBray	44.71	−0.77	62	900	13.0	Forest	L. Wingate & J. Ogée, unpublished
France_Barbeta	Ciron	44.38	−0.31	60	813	12.9	Forest	A. Barbeta, unpublished
Canada_Flanagan	Lethbridge	49.69	−112.83	910	380	5.8	Grassland	(Flanagan <i>et al.</i> , 1991a)

temperature, then w_a/w_i is equal to the relative humidity of the air surrounding the leaf. Eqn 1 is an approximation of a more precise form of the Craig–Gordon equation (Farquhar *et al.*, 2007); however, it is very useful in that it shows intuitively what the different drivers of δ_e are expected to be. Therefore, we used the more precise version of the equation for calculations and analyses, but used the approximate version here to guide our discussion. A summary of formulae for calculating the equilibrium and kinetic fractionation factors for $\delta^{18}\text{O}$ and $\delta^2\text{H}$ and the more precise version of the Craig–Gordon equation can be found in Cernusak *et al.* (2016) and in the Supporting Information Dataset S1.

Eqn 1 assumes isotopic steady state, in which the water leaving the leaf through transpiration has the same isotopic composition as that entering the leaf from the xylem. Furthermore, it makes a prediction for the evaporative sites, while the unit of measure in our dataset is bulk leaf water (δ_l), the total sum of water extracted from the leaf. Bulk leaf water can be expected to be somewhat less enriched than the evaporative sites, due to the influx of unenriched xylem water in the veins (Roden & Ehleringer, 1999; Farquhar & Gan, 2003; Farquhar *et al.*, 2007; Holloway-Phillips *et al.*, 2016). Whereas the mechanisms in the Craig–Gordon equation are

identical for $\delta^2\text{H}$ and $\delta^{18}\text{O}$, the relative magnitudes of the equilibrium and kinetic fractionation factors differ. For $\delta^2\text{H}$, the ϵ^+ is relatively large and ϵ_k relatively small, whereas the converse is true for $\delta^{18}\text{O}$ (Merlivat, 1978; Horita & Wesolowski, 1994; Cernusak *et al.*, 2016). The ratio $\epsilon^+ : \epsilon_k$ is *c.* 3 : 1 for $\delta^2\text{H}$ and 1 : 3 for $\delta^{18}\text{O}$ (Dataset S1).

We plotted the Craig–Gordon predicted leaf water isotopic compositions against observations for our dataset, to determine whether the Craig–Gordon equation could provide a reasonable framework for guiding analyses of different drivers. Fig. 1 shows the observed bulk leaf water $\delta^2\text{H}$ and $\delta^{18}\text{O}$ plotted against that predicted by the Craig–Gordon equation, using the measured air temperature, relative humidity, isotopic composition of xylem water and atmospheric vapour. Overall, the Craig–Gordon equation explained 89% of observed variation in leaf water $\delta^2\text{H}$ and 67% of observed variation in leaf water $\delta^{18}\text{O}$. As anticipated, the slopes of the relationships were less than unity, as would be the case if some fraction of bulk leaf water represented unenriched xylem water. The generally good predictive ability of the Craig–Gordon equation for daytime leaf water isotopic composition suggests that it can provide a framework for evaluating whether




Fig. 1 Observed leaf water isotopic composition for $\delta^2\text{H}$ (a) and $\delta^{18}\text{O}$ (b) plotted against values predicted by the Craig–Gordon equation. Symbols with different colours refer to the different datasets that have been compiled for this paper. The dotted lines show one-to-one lines and solid lines show least-squares linear regressions with fitted coefficients shown in the panels along with the coefficient of determination, R^2 .

different environmental drivers predominate for hydrogen vs oxygen.

Some additional sources of unexplained variation in Fig. 1 could include departures from isotopic steady state (Dongmann *et al.*, 1974; Farquhar & Cernusak, 2005), variation in the fraction of unenriched water in leaves associated with differences in leaf anatomy and physiology (Holloway-Phillips *et al.*, 2016; Barbour *et al.*, 2021) and unaccounted for variation in boundary layer conductance (Buhay *et al.*, 1996). The detailed data required to test for each of these possibilities were not available across the compiled dataset. However, we did repeat our analyses with observations limited to the middle of the day (from 11:00 h to 14:00 h), when isotopic steady state is most likely to be achieved (Harwood *et al.*, 1998). This yielded very similar results to those shown in Fig. 1. The same was also true for subsequent figures and we therefore present analyses with all daytime observations included.

The environmental drivers that are used in the Craig–Gordon equation are air temperature, which impacts ε^+ (Horita &

Wesolowski, 1994); relative humidity, which is assumed equal to w_a/w_l if leaf temperature has not deviated from air temperature and w_l is saturated; isotopic composition of source water entering the leaf, assumed equal to the measured xylem water in our analyses; and the isotopic composition of atmospheric water vapour. Fig. 2 shows the observed leaf water $\delta^2\text{H}$ and $\delta^{18}\text{O}$ plotted against each of these four environmental drivers. For $\delta^2\text{H}_l$, xylem water $\delta^2\text{H}$ and atmospheric vapour $\delta^2\text{H}$ were much more strongly correlated with it than air temperature or relative humidity. For $\delta^{18}\text{O}_l$, conversely, air relative humidity was much more strongly correlated than any of the other drivers. For $\delta^2\text{H}_l$, either xylem water or atmospheric vapour $\delta^2\text{H}$ explained more than two-thirds of its variation, whereas for $\delta^{18}\text{O}_l$ the air relative humidity explained about half of its variation.

The reason that xylem water is a much stronger driver of leaf water for $\delta^2\text{H}$ than for $\delta^{18}\text{O}$ is because the range of variation in meteoric water isotopic composition compared with that in leaf water evaporative enrichment is larger for $\delta^2\text{H}$ than for $\delta^{18}\text{O}$. This





Fig. 2 Observed isotopic composition for leaf water $\delta^2\text{H}$ (a–d) and $\delta^{18}\text{O}$ (e–h) plotted against the four environmental drivers in the Craig–Gordon equation: air temperature (a, e), air relative humidity (b, f), the corresponding isotopic composition of atmospheric vapour (c, g) and the corresponding isotopic composition of xylem water (d, h). The symbol colours show the different datasets compiled for this paper. Solid lines are least-squares linear regressions, with fitted coefficients shown in the panels, along with the coefficient of determination, R^2 .

Fig. 3 The isotopic composition of xylem water, leaf water and atmospheric vapour plotted in $\delta^2\text{H}$ – $\delta^{18}\text{O}$ dual-isotope space. The black line shows the meteoric water line, defined as $\delta^2\text{H} = 8 \times \delta^{18}\text{O} + 10$. The coloured lines show the evaporation lines for leaf water, in which the intercept with the meteoric water line is the mean for each site and the slope is calculated as $(\delta^2\text{H}_l - \delta^2\text{H}_x) / (\delta^{18}\text{O}_l - \delta^{18}\text{O}_x)$ using the mean quantities for each site, where subscript 'l' refers to leaf water and 'x' to xylem water. The colours of the lines refer to the individual datasets compiled for this paper. The range of $\delta^{18}\text{O}_l$ observed for each site defines the length of the coloured lines.



can be seen through inspection of Fig. 3, which shows the evaporation lines for leaf water for each site in the dataset and their extrapolation to the meteoric water line. The range in the y -axis over which the evaporation lines intersect the meteoric water line for $\delta^2\text{H}$ is $c. 120\text{‰}$ and the corresponding range on the x -axis for $\delta^{18}\text{O}$ is $c. 15\text{‰}$, for a ratio of $c. 8 : 1$, consistent with the slope of the meteoric water line. Conversely, the range for leaf water isotopic composition beginning at the meteoric water line and moving right along the evaporation lines is $c. 100\text{‰}$ on the y -axis for $\delta^2\text{H}$ and $c. 30\text{‰}$ on the x -axis for $\delta^{18}\text{O}$, for a ratio of $c. 3 : 1$. Therefore the point at which the evaporation line intersects the meteoric water line can exert a much stronger influence on leaf water for $\delta^2\text{H}$ than for $\delta^{18}\text{O}$, because its range is relatively large compared with the range over which evaporation can enrich the leaf water above source water. Another way to understand this conceptually is to consider that the slope of the meteoric water line, defining source water variation in $\delta^2\text{H}$ – $\delta^{18}\text{O}$ space, corresponds approximately to the ratio of the equilibrium fractionations for $\delta^2\text{H}$ and $\delta^{18}\text{O}$ (mean = 8.6 in our dataset). Conversely, the slopes of the evaporation lines corresponded approximately to the ratio of the sum of equilibrium and kinetic fractionations (mean = 2.9 in our dataset).

This difference between leaf water dynamics for $\delta^2\text{H}$ and $\delta^{18}\text{O}$, driven by source water isotopic composition, is important for interpreting organic material signals. For example, leaf water proxies based on $\delta^2\text{H}$, such as the $\delta^2\text{H}$ of n -alkanes derived from leaf waxes, if sampled across a large geographic range, could be expected to be strongly influenced by a widely varying $\delta^2\text{H}$ of source water (Liu & Yang, 2008; Sachse *et al.*, 2012; Ladd *et al.*, 2021). Conversely, only if there is little variation in source water $\delta^2\text{H}$, will the variation in n -alkane $\delta^2\text{H}$ of leaf waxes reliably record the extent of leaf water evaporative enrichment (Kahmen *et al.*, 2013b). For an organic matter proxy such as cellulose $\delta^{18}\text{O}$, we would expect the geographic variation in source water isotopic composition to have less influence compared with the dynamics of

leaf water enrichment above source water, driven primarily by relative humidity (Barbour & Farquhar, 2000; Kahmen *et al.*, 2011). To the extent that such geographic variation can provide a space for time substitution, our results also have implications for interpreting changes through time within a site. For example, $\delta^2\text{H}$ of n -alkanes from leaf waxes has been combined with $\delta^{18}\text{O}$ of hemicellulose sugars for reconstructing paleoclimate from sedimentary records (Zech *et al.*, 2013; Hepp *et al.*, 2021). Our results suggest that $\delta^2\text{H}$ of n -alkanes should be better suited to detecting changes in $\delta^2\text{H}$ of precipitation and $\delta^{18}\text{O}$ of hemicellulose sugars to detecting changes in relative humidity. We note, however, that the extent of transfer of the leaf water signal to the biomarker will also be important; for example, for cellulose $\delta^{18}\text{O}$, our analysis of leaf water $\delta^{18}\text{O}$ may be more relevant to leaf cellulose than to stem wood cellulose, as the latter is subject to partial exchange with unenriched xylem water (Roden *et al.*, 2000; Kahmen *et al.*, 2011; Cheesman & Cernusak, 2017), with the same caveat also applicable for grasses (Helliker & Ehleringer, 2000; Liu *et al.*, 2017).

Isotopic enrichment of leaf water above xylem water

In addition to xylem water $\delta^2\text{H}$ having a close correlation with leaf water $\delta^2\text{H}$, atmospheric vapour $\delta^2\text{H}$ also had a close correlation. The next question we asked in our analysis was whether the relationship between leaf water and vapour for $\delta^2\text{H}$ would still remain stronger than that for $\delta^{18}\text{O}$ when variation in xylem water isotopic composition was removed. To answer this question, we expressed leaf water as enrichment above xylem water (Δ_l), calculated as $\Delta_l = (\delta_l - \delta_x) / (1 + \delta_x)$, where the subscript 'l' refers to leaf water and 'x' to xylem water. Again, we use the approximate form of the Craig–Gordon equation here for ease of interpretation to guide our analysis, but used the more precise form in our calculations. With leaf water expressed as enrichment above source water, the Craig–Gordon equation becomes (Farquhar *et al.*, 1989):




Fig. 4 Leaf water isotopic enrichment above xylem water for hydrogen, $\Delta^2\text{H}$ (a–c), and for oxygen, $\Delta^{18}\text{O}$ (d–f), plotted against air temperature (a, d), air relative humidity (b, e) and the corresponding isotopic enrichment of atmospheric vapour above xylem water (c, f). Solid lines show least-squares linear regressions, along with fitted coefficients and the coefficient of determination, R^2 . Symbol colours refer to individual datasets compiled for this paper.

$$\Delta_e \approx \epsilon^+ + \epsilon_k + (\Delta_v - \epsilon_k)h$$

Eqn 2

(Δ_e , predicted enrichment at the evaporative sites in leaves and Δ_v , enrichment of atmospheric vapour relative to source water). Note that this latter term is generally negative; that is, atmospheric vapour is generally depleted in heavier isotopes compared with source water. In our analysis, we calculated Δ_v as $\Delta_v = (\delta_v - \delta_x) / (1 + \delta_x)$, where δ_v is $\delta^2\text{H}$ or $\delta^{18}\text{O}$ of atmospheric vapour and δ_x is that of xylem water. In Fig. 4, we show the observed bulk leaf water enrichment plotted against the three environmental drivers remaining in Eqn 2.

Fig. 4 shows that the correlation between leaf water and relative humidity for hydrogen has been markedly improved by removing source water variation, with relative humidity now explaining 41% of the variation in $\Delta^2\text{H}_l$. Therefore, after calculating $\Delta^2\text{H}_l$ to remove the source water signal, the sensitivity to relative humidity became more apparent. For oxygen, there was also a strengthening of the correlation between leaf water enrichment and relative humidity, with the R^2 increasing from 0.49 to 0.57. Interestingly, however, the correlation between leaf water enrichment and atmospheric vapour enrichment was still relatively strong for hydrogen, but weak for oxygen, which stands out as a point of difference between $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ in Fig. 4. Stronger relationships with atmospheric vapour for $\Delta^2\text{H}_l$ than for $\Delta^{18}\text{O}_l$ have also been observed previously in some of the individual datasets that have now been compiled for this paper (Cernusak *et al.*, 2016; Bögelein *et al.*, 2017; Munksgaard *et al.*, 2017).

Role of atmospheric vapour isotopic composition

Can we identify further the underlying cause of the stronger correlation between leaf water and atmospheric vapour for $\Delta^2\text{H}$ compared with $\Delta^{18}\text{O}$? To explore this, we turned again to the Craig–Gordon equation, taking the derivative of Eqn 2 with respect to Δ_v . This provides a mathematical description of predicted drivers of the change in Δ_l for a given change in Δ_v :

$$\frac{d\Delta_l}{d\Delta_v} = \frac{d\epsilon^+}{d\Delta_v} + \frac{d\epsilon_k}{d\Delta_v} + h \left(1 - \frac{d\epsilon_k}{d\Delta_v} \right) + (\Delta_v - \epsilon_k) \frac{dh}{d\Delta_v} \quad \text{Eqn 3}$$

We used our dataset to estimate the terms in Eqn 3 by taking regression slopes for the derivative terms and mean values for h and $(\Delta_v - \epsilon_k)$. These estimates are shown in Table 2 for $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$. From Table 2, it can be seen that the first two terms on the right side of Eqn 3 are small in magnitude for both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ and unlikely to have a strong influence on $d\Delta_l/d\Delta_v$ for either. Because $d\epsilon_k/d\Delta_v$ is small, it means that the third term on the right side will approach the value of h , which is larger by comparison, having a mean value in our dataset of 0.5, or an air relative humidity of *c.* 50%. The largest term in Eqn 3 by far for both $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$ is $(\Delta_v - \epsilon_k)$, having mean values of -79‰ and -35‰ for $\Delta^2\text{H}$ and $\Delta^{18}\text{O}$, respectively. This is then multiplied by a much smaller term, $dh/d\Delta_v$. Importantly, $(\Delta_v - \epsilon_k)$ is negative, setting up the possibility that the interplay between the third and fourth terms on the right side of the equation could be important, with the third term, $h(1 - d\epsilon_k/d\Delta_v)$, being positive and the fourth term,

Table 2 Values for the terms in Eqn 3 calculated from the combined dataset.

	$\frac{d\epsilon_v}{d\Delta_v}$	$\frac{d\epsilon_k}{d\Delta_v}$	h	$(\Delta_v - \epsilon_k)$	$\frac{dh}{d\Delta_v}$	$h\left(1 - \frac{d\epsilon_k}{d\Delta_v}\right)$	$(\Delta_v - \epsilon_k) \frac{dh}{d\Delta_v}$	Sum of shaded columns
$\Delta^2\text{H}$	−0.09	−0.01	0.51	−79.2	−0.0004	0.51	0.03	0.45
$\Delta^{18}\text{O}$	0.01	−0.02	0.51	−35.2	0.0102	0.52	−0.36	0.15

Derivative terms (dy/dx) were calculated as the slope of a linear regression of the two parameters y and x , whereas nonderivative terms were calculated as the mean of the given parameter. According to Eqn 3, the dependence of leaf water isotopic enrichment on the atmospheric vapour isotopic composition, $d\Delta_l/d\Delta_v$, is equal to the sum of the shaded columns, which is shown in the final column. As can be seen, the primary difference for $\Delta^2\text{H}$ compared with $\Delta^{18}\text{O}$ results from the term $dh/d\Delta_v$; that is, a correlation between atmospheric humidity and the isotopic composition of atmospheric vapour, which is much stronger for $\Delta^{18}\text{O}$ than for $\Delta^2\text{H}$.

$(\Delta_v - \epsilon_k)d h/d\Delta_v$, potentially counteracting it with a negative value.

The interaction between the third and fourth terms in Eqn 3 does indeed appear to be pivotal in explaining why leaf water correlates more strongly with atmospheric vapour for $\Delta^2\text{H}$ than for $\Delta^{18}\text{O}$. For $\Delta^2\text{H}$, the linear regression between h and $\Delta^2\text{H}_v$ was not significant ($P = 0.32$, $n = 546$) and had a slope of -0.0004 . This gives a value for the fourth term in Eqn 3 for $\Delta^2\text{H}$ of 0.03, which therefore adds slightly to the positive value of the third term, again having a value of $c. 0.5$. Conversely, for $\Delta^{18}\text{O}$, the regression between h and $\Delta^{18}\text{O}_v$ was significant ($P < 0.001$, $n = 546$) and had a positive slope of 0.0102. Because this slope was positive, the fourth term on the right side of Eqn 3 for $\Delta^{18}\text{O}$ takes on an overall negative value of -0.36 . Therefore, for $\Delta^{18}\text{O}$, the fourth term on the right side of Eqn 3 largely cancels the influence of the third term and, as a result, there is little change in $\Delta^{18}\text{O}_l$ for a given change in $\Delta^{18}\text{O}_v$. This manifests in our dataset as a weakened correlation between leaf water and atmospheric vapour for $\Delta^{18}\text{O}$ as seen in Fig. 4f whereas, for $\Delta^2\text{H}$, there is a stronger correlation between leaf water and atmospheric vapour, as seen in Fig. 4c.

The analysis above shows that there is a positive correlation between relative humidity and the $\Delta^{18}\text{O}$ of atmospheric vapour in our dataset, whereas such a correlation does not exist between relative humidity and the $\Delta^2\text{H}$ of atmospheric vapour. Through application of Eqn 3, we showed that this difference partly explains why leaf water $\delta^2\text{H}$ more strongly correlates with atmospheric vapour $\delta^2\text{H}$ than is the case for $\delta^{18}\text{O}$. Another way to approach the underlying issue of this apparent difference in behaviour of atmospheric vapour for the two isotopologues with respect to relative humidity is to calculate the deuterium excess or the departure from an expectation of the relationship between $\delta^2\text{H}$ and $\delta^{18}\text{O}$ based on the meteoric water line. The global meteoric water line is described by $\delta^2\text{H} = 8 \times \delta^{18}\text{O} + 10$ (Craig, 1961). We therefore calculated the deuterium excess of water vapour, d_v , as $d_v = \delta^2\text{H}_v - 8 \times \delta^{18}\text{O}_v$ (Dansgaard, 1964). We also made the same calculation for atmospheric vapour composition with respect to xylem water, $\Delta_{dv} = \Delta^2\text{H}_v - 8 \times \Delta^{18}\text{O}_v$. We then tested for correlations between these parameters and the air relative humidity in our dataset. Both showed significant negative correlations with relative humidity, with the relationship stronger for Δ_{dv} ($R^2 = 0.17$, $P < 0.001$, $n = 546$) than for d_v ($R^2 = 0.08$, $P < 0.001$, $n = 546$).

A relationship between the deuterium excess of atmospheric vapour and relative humidity has also been observed on diurnal timescales at six sites in the northern hemisphere (Welp *et al.*, 2012) and at a tropical site in Cairns, Australia (Munksgaard *et al.*, 2020). This pronounced, general pattern is thought to be driven by the diurnal pattern of plant transpiration and the contribution of transpired water to atmospheric vapour and by entrainment of the free atmosphere into the planetary boundary layer with increased convective mixing during the day. The result is a general midday decrease in the $\delta^{18}\text{O}$ of atmospheric vapour, but little to no change in $\delta^2\text{H}$. This leads to the diurnal variability of d_v , which is anticorrelated with the diurnal pattern of relative humidity (Welp *et al.*, 2012; Munksgaard *et al.*, 2020). A shorter time series of 3 d at the Wind River Experimental Forest (Washington, USA) showed a similar pattern (Lai & Ehleringer, 2011). The strength of this pattern suggests that such diurnal variation could be driving the overall relationship between d_v and relative humidity in our dataset. Welp *et al.* (2012) also observed negative correlations between day-to-day variation in d_v and relative humidity throughout the summer months at sites located near large bodies of water, with such patterns also previously reported for sites in marine-type settings (Uemura *et al.*, 2008; Gat *et al.*, 2011). Such dynamics related to marine air sources may also have been relevant at some sites within our dataset. When we restricted our analysis from daytime observations to only midday observations (between 11:00 h and 13:00 h) to minimise diurnality, we observed a weak, but still significant, relationship between d_v and relative humidity ($R^2 = 0.02$, $P < 0.05$, $n = 200$), showing the importance of diurnal effects. What is clear overall is that covariation between $\Delta^{18}\text{O}_v$ and relative humidity, but not $\Delta^2\text{H}_v$ and relative humidity, plays an important role in modulating leaf water isotope dynamics, leading to the result highlighted in our dataset of $\Delta^2\text{H}$ of leaf water showing stronger correlation with $\Delta^2\text{H}$ of atmospheric vapour than is the case for $\Delta^{18}\text{O}$.

Are two isotopes better than one?

Some organic matter proxies, such as plant cellulose, allow both $\delta^{18}\text{O}$ and $\delta^2\text{H}$ to be measured. In such cases, it is possible to estimate leaf water isotopic composition for both $\delta^2\text{H}$ and $\delta^{18}\text{O}$ and therefore to reconstruct d_l , the deuterium excess for leaf water. As seen in Fig. 3, the slopes of the evaporation lines tend to be uniform and independent of the source water and the position




Fig. 5 Deuterium excess in leaf water plotted against (a) air temperature, (b) air relative humidity, (c) the deuterium excess in atmospheric water vapour and (d) the deuterium excess in xylem water. The deuterium excess, d , was calculated as $d = \delta^2\text{H} - 8 \times \delta^{18}\text{O}$. Solid lines show least-squares linear regressions, along with fitted coefficients and the coefficient of determination, R^2 . Symbol colours refer to individual datasets compiled for this paper.

along the evaporation line is determined mainly by relative humidity. Therefore, d_1 is expected to show a relationship with relative humidity that is largely independent of source water isotope composition. For this reason, it has been suggested that such a dual-isotope approach could provide a stronger basis for reconstructing relative humidity than either isotope alone when the source water isotopic composition is not known (Zech *et al.*, 2013; Voelker *et al.*, 2014). Our dataset gave us an opportunity to test this idea across a diverse range of sites and conditions.

Fig. 5 shows d_1 for the dataset, calculated as $d_1 = \delta^2\text{H}_1 - 8 \times \delta^{18}\text{O}_1$, plotted against potential drivers, including relative humidity. Because d_1 was calculated from $\delta^{18}\text{O}_1$ and $\delta^2\text{H}_1$, we can compare the relationships with relative humidity among the three. The d_1 does indeed have the strongest relationship; however, it is not very much stronger than that for $\delta^{18}\text{O}_1$. The R^2 for d_1 vs relative humidity is 0.54 (Fig. 5b), whereas that for $\delta^{18}\text{O}_1$ is 0.49 (Fig. 2f), and that for $\delta^2\text{H}_1$ is 0.07 (Fig. 2b). Therefore, surprisingly, leaf water $\delta^{18}\text{O}$ on its own would be nearly as good a predictor of air relative humidity as d_1 calculated from both the $\delta^{18}\text{O}$ and $\delta^2\text{H}$ of leaf water. The explanation for this is the relatively constrained variation in $\delta^{18}\text{O}$ of xylem water, especially in comparison with $\delta^2\text{H}$ (Fig. 3) and the relative insensitivity of $\Delta^{18}\text{O}_1$ to atmospheric vapour $\Delta^{18}\text{O}_v$, as discussed above. Given these considerations, we suggest that if applying a dual-isotope approach in this way, one should weigh up carefully the uncertainty associated with estimating leaf water $\delta^2\text{H}$ from cellulose $\delta^2\text{H}$, given the relative complexity in the signal transfer pathway from leaf water to cellulose for $\delta^2\text{H}$ (Cormier *et al.*, 2018; Lehmann *et al.*, 2021). The cost in uncertainty associated with this may not be worth the relatively modest improvement in strength of the correlation between d_1 and relative humidity compared with that for $\delta^{18}\text{O}$ alone.

Conclusions

We compiled a dataset for $\delta^2\text{H}$ and $\delta^{18}\text{O}$ of leaf water, xylem water, atmospheric vapour, air temperature and relative humidity from a diverse range of sites and used the dataset to test for differences in how leaf water $\delta^2\text{H}$ and $\delta^{18}\text{O}$ reflect environmental drivers. We conducted the analysis in the context of asking which drivers could be best reconstructed from leaf water proxies based on organic

material. Xylem water $\delta^2\text{H}$ was a much stronger driver of variation in leaf water $\delta^2\text{H}$ than was the case for xylem water $\delta^{18}\text{O}$ as a driver of variation in leaf water $\delta^{18}\text{O}$. Conversely, relative humidity showed a considerably stronger relationship with leaf water $\delta^{18}\text{O}$ than it did with leaf water $\delta^2\text{H}$. This pattern persisted when we removed xylem water isotopic variation from the leaf water signal by expressing the leaf water isotopic composition as an enrichment above xylem water. We identified the underlying reason for this pattern as a correlation between relative humidity and the $\delta^{18}\text{O}$ of atmospheric vapour. Such a correlation has also been observed in time series of vapour isotopic composition measurements, and manifests most clearly as an anticorrelation between the deuterium excess of atmospheric vapour and relative humidity on diurnal timescales. While we did not have sufficient resolution of sampling within sites to tease this apart in our dataset, we suspect that this diurnal pattern is likely to underlie the correlation between relative humidity and atmospheric vapour $\Delta^{18}\text{O}_v$ that we observed. We conclude that leaf water $\delta^2\text{H}$ and $\delta^{18}\text{O}$ do indeed reflect the balance of potential environmental drivers differently: leaf water $\delta^2\text{H}$ reflects more strongly xylem water $\delta^2\text{H}$ and atmospheric vapour $\delta^2\text{H}$, whereas leaf water $\delta^{18}\text{O}$ reflects more strongly air relative humidity.

Acknowledgements

LAC acknowledges the support of a Make Our Planet Great Again Fellowship through Campus France to visit MC at INRAE, UMR Silva, Nancy to work on this project. MC was supported by a grant overseen by the French National Research Agency (ANR) as part of the 'Investissements d'Avenir' programme (ANR-11-LABX-0002-01, Laboratory of Excellence ARBRE) and by a grant from the German Research Foundation (DFG, CU 173/4-1). L. Wang acknowledges partial support from the Division of Earth Sciences of the US National Science Foundation (NSF EAR-1554894).

Author contributions

LAC and MC initiated the review and conducted initial analyses. All authors contributed to further development of ideas and writing of the manuscript.

ORCID

Adrià Barbeta  <https://orcid.org/0000-0002-8357-1719>
 Lucas A. Cernusak  <https://orcid.org/0000-0002-7575-5526>
 Matthias Cuntz  <https://orcid.org/0000-0002-5966-1829>
 Rebekka Eichstaedt (Bögelein)  <https://orcid.org/0000-0002-4905-7994>
 Juan Pedro Ferrio  <https://orcid.org/0000-0001-5904-7821>
 Lawrence B. Flanagan  <https://orcid.org/0000-0003-1748-0306>
 Arthur Gessler  <https://orcid.org/0000-0002-1910-9589>
 Ansgar Kahmen  <https://orcid.org/0000-0002-7823-5163>
 Claudia Keitel  <https://orcid.org/0000-0003-3101-7529>
 Chun-Ta Lai  <https://orcid.org/0000-0001-9662-8545>
 Paula Martín-Gómez  <https://orcid.org/0000-0001-5810-6762>
 Niels C. Munksgaard  <https://orcid.org/0000-0003-4906-1544>
 Daniel B. Nelson  <https://orcid.org/0000-0002-2716-7770>
 Jérôme Ogée  <https://orcid.org/0000-0002-3365-8584>
 John S. Roden  <https://orcid.org/0000-0001-8641-1625>
 Hans Schnyder  <https://orcid.org/0000-0002-0139-7535>
 Hilary Stuart-Williams  <https://orcid.org/0000-0002-9272-9178>
 Steven L. Voelker  <https://orcid.org/0000-0002-0110-3381>
 Lixin Wang  <https://orcid.org/0000-0003-0968-1247>
 Lisa Wingate  <https://orcid.org/0000-0003-1921-1556>
 Wusheng Yu  <https://orcid.org/0000-0002-3089-0820>
 Liangju Zhao  <https://orcid.org/0000-0001-7445-3916>

Data availability

All data used in the analyses presented in this paper are available in Supporting Information Dataset S1.

References

- Ayliffe LK, Chivas AR. 1990. Oxygen isotope composition of the bone phosphate of Australian kangaroos: potential as a palaeoenvironmental recorder. *Geochimica et Cosmochimica Acta* 54: 2603–2609.
- Barbata A, Burrett R, Martín-Gómez P, Fréjaville B, Devert N, Wingate L, Domec J-C, Ogée J. 2022. Evidence for distinct isotopic compositions of sap and tissue water in tree stems: consequences for plant water source identification. *New Phytologist* 233: 1121–1132.
- Barbour MM. 2007. Stable oxygen isotope composition of plant tissue: a review. *Functional Plant Biology* 34: 83–94.
- Barbour MM, Farquhar GD. 2000. Relative humidity- and ABA-induced variation in carbon and oxygen isotope ratios of cotton leaves. *Plant, Cell & Environment* 23: 473–485.
- Barbour MM, Farquhar GD, Buckley TN. 2017. Leaf water stable isotopes and water transport outside the xylem. *Plant, Cell & Environment* 40: 914–920.
- Barbour MM, Loucos KE, Lockhart EL, Shrestha A, McCallum D, Simonin KA, Song X, Griffani DS, Farquhar GD. 2021. Can hydraulic design explain patterns of leaf water isotopic enrichment in C₃ plants? *Plant, Cell & Environment* 44: 432–444.
- Bögelein R, Thomas FM, Kahmen A. 2017. Leaf water ¹⁸O and ²H enrichment along vertical canopy profiles in a broadleaved and a conifer forest tree. *Plant, Cell & Environment* 40: 1086–1103.
- Buckley TN, Sack L. 2019. The humidity inside leaves and why you should care: implications of unsaturation of leaf intercellular airspaces. *American Journal of Botany* 106: 618–621.
- Buhay WM, Edwards TWD, Aravena R. 1996. Evaluating kinetic fractionation factors used for ecologic and paleoclimatic reconstructions from oxygen and hydrogen isotope ratios in plant water and cellulose. *Geochimica et Cosmochimica Acta* 60: 2209–2218.
- Bush RT, Berke MA, Jacobson AD. 2017. Plant water δD and δ¹⁸O of tundra species from west Greenland. *Arctic, Antarctic, and Alpine Research* 49: 318.
- Cernusak LA, Barbour MM, Arndt SK, Cheesman AW, English NB, Feild TS, Helliker BR, Holloway-Phillips MM, Holtum JAM, Kahmen A *et al.* 2016. Stable isotopes in leaf water of terrestrial plants. *Plant, Cell & Environment* 39: 1087–1102.
- Cernusak LA, Farquhar GD, Pate J. 2005. Environmental and physiological controls over oxygen and carbon isotope composition of Tasmanian blue gum, *Eucalyptus globulus*. *Tree Physiology* 25: 129–146.
- Cernusak LA, Goldsmith G, Arend M, Siegwolf RTW. 2019. Effect of vapor pressure deficit on gas exchange in wild-type and abscisic acid-insensitive plants. *Plant Physiology* 181: 1573–1586.
- Cernusak LA, Pate JS, Farquhar GD. 2002. Diurnal variation in the stable isotope composition of water and dry matter in fruiting *Lupinus angustifolius* under field conditions. *Plant, Cell & Environment* 25: 893–907.
- Cernusak LA, Ubierna N, Jenkins MW, Garrity SR, Rahn T, Powers HH, Hanson DT, Sevanto S, Wong SC, McDowell NG *et al.* 2018. Unsaturation of vapour pressure inside leaves of two conifer species. *Scientific Reports* 8: 7667.
- Cheesman AW, Cernusak LA. 2017. Infidelity in the outback: climate signal recorded in Δ¹⁸O of leaf but not branch cellulose of eucalypts across an Australian aridity gradient. *Tree Physiology* 37: 554–564.
- Chen Y, Helliker BR, Tang X, Li F, Zhou Y, Song X. 2020. Stem water cryogenic extraction biases estimation in deuterium isotope composition of plant source water. *Proceedings of the National Academy of Sciences, USA* 117: 33345.
- Cormier M-A, Werner RA, Sauer PE, Gröcke DR, Leuenberger MC, Wieloch T, Schleucher J, Kahmen A. 2018. ²H-fractionations during the biosynthesis of carbohydrates and lipids imprint a metabolic signal on the δ²H values of plant organic compounds. *New Phytologist* 218: 479–491.
- Craig H. 1961. Isotopic variations in meteoric waters. *Science* 133: 1702–1703.
- Craig H, Gordon LI. 1965. Deuterium and oxygen-18 variations in the ocean and the marine atmosphere. In: Tongiorgi E, ed. *Proceedings of a conference on stable isotopes in oceanographic studies and palaeotemperatures*. Pisa, Italy: Lisch and Figli, 9–130.
- Cuntz M, Ogée J, Farquhar GD, Peylin P, Cernusak LA. 2007. Modelling advection and diffusion of water isotopologues in leaves. *Plant, Cell & Environment* 30: 892–909.
- Dansgaard W. 1964. Stable isotopes in precipitation. *Tellus* 16: 436–468.
- Dongmann G, Nurnberg HW, Förstel H, Wagener K. 1974. On the enrichment of H₂¹⁸O in the leaves of transpiring plants. *Radiation and Environmental Biophysics* 11: 41–52.
- Faith JT. 2018. Paleodietary change and its implications for aridity indices derived from δ¹⁸O of herbivore tooth enamel. *Palaeogeography, Palaeoclimatology, Palaeoecology* 490: 571–578.
- Farquhar GD, Barbour MM, Henry BK. 1998. Interpretation of oxygen isotope composition of leaf material. In: Griffiths H, ed. *Stable isotopes: integration of biological, ecological, and geochemical processes*. Oxford, UK: BIOS Scientific, 27–48.
- Farquhar GD, Cernusak LA. 2005. On the isotopic composition of leaf water in the non-steady state. *Functional Plant Biology* 32: 293–303.
- Farquhar GD, Cernusak LA, Barnes B. 2007. Heavy water fractionation during transpiration. *Plant Physiology* 143: 11–18.
- Farquhar GD, Gan KS. 2003. On the progressive enrichment of the oxygen isotopic composition of water along leaves. *Plant, Cell & Environment* 26: 801–819.
- Farquhar GD, Hubick KT, Condon AG, Richards RA. 1989. Carbon isotope fractionation and plant water-use efficiency. In: Rundel PW, Ehleringer JR, Nagy KA, eds. *Stable isotopes in ecological research*. New York, NY, USA: Springer-Verlag, 21–46.
- Flanagan LB, Bain JF, Ehleringer JR. 1991a. Stable oxygen and hydrogen isotope composition of leaf water in C₃ and C₄ plant species under field conditions. *Oecologia* 88: 394–400.
- Flanagan LB, Comstock JP, Ehleringer JR. 1991b. Comparison of modeled and observed environmental influences on the stable oxygen and hydrogen isotope composition of leaf water in *Phaseolus vulgaris* L. *Plant Physiology* 96: 588–596.
- Flanagan LB, Marshall JD, Ehleringer JR. 1993. Photosynthetic gas exchange and the stable isotope composition of leaf water - comparison of a xylem-tapping mistletoe and its host. *Plant, Cell & Environment* 16: 623–631.

- Gan KS, Wong SC, Yong JWH, Farquhar GD. 2002. ^{18}O spatial patterns of vein xylem water, leaf water, and dry matter in cotton leaves. *Plant Physiology* 130: 1008–1021.
- Gat JR, Klein B, Kushnir Y, Roether W, Wernli H, Yam R, Shemesh A. 2011. Isotope composition of air moisture over the Mediterranean sea: an index of the air-sea interaction pattern. *Tellus B: Chemical and Physical Meteorology* 55: 953–965.
- Goldsmith GR, Lehmann MM, Cernusak LA, Arend M, Siegwolf RTW. 2017. Inferring foliar water uptake using stable isotopes of water. *Oecologia* 184: 763–766.
- Gonfiantini R, Gratzu S, Tongiorgi E. 1965. Oxygen isotope composition of water in leaves. In: International Atomic Energy Agency, ed. *Isotopes and radiation in soil-plant nutrition studies*. Vienna, Austria: IAEA, 405–410.
- Harwood KG, Gillon JS, Griffiths H, Broadmeadow MSJ. 1998. Diurnal variation of $\Delta^{13}\text{CO}_2$, $\Delta^{18}\text{O}^{16}\text{O}$, and evaporative site enrichment of $\delta\text{H}_2^{18}\text{O}$ in *Piper aduncum* under field conditions in Trinidad. *Plant, Cell & Environment* 21: 269–283.
- Helliker BR. 2014. Reconstructing the $\delta^{18}\text{O}$ of atmospheric water vapour via the CAM epiphyte *Tillandsia usneoides*: seasonal controls on $\delta^{18}\text{O}$ in the field and large-scale reconstruction of $\delta^{18}\text{O}_{\text{a}}$. *Plant, Cell & Environment* 37: 541–556.
- Helliker BR, Ehleringer JR. 2000. Establishing a grassland signature in veins: ^{18}O in the leaf water of C_3 and C_4 grasses. *Proceedings of the National Academy of Sciences, USA* 97: 7894–7898.
- Hepp J, Mayr C, Rozanski K, Schäfer IK, Tuthorn M, Glaser B, Juchelka D, Stichter W, Zech R, Zech M. 2021. Validation of a coupled $\delta^2\text{H}_{\text{n-alkane}}-\delta^{18}\text{O}_{\text{sugar}}$ paleohygrometer approach based on a climate chamber experiment. *Biogeosciences* 18: 5363–5380.
- Hirt RL, Schnyder H, Ostler U, Schäufele R, Schleip I, Vetter SH, Auerswald K, Baca Cabrera JC, Wingate L, Barbour MM *et al.* 2019. The ^{18}O ecohydrology of a grassland ecosystem – predictions and observations. *Hydrology and Earth System Sciences* 23: 2581–2600.
- Holloway-Phillips M-M, Cernusak LA, Barbour M, Song X, Cheesman A, Munksgaard N, Stuart-Williams H, Farquhar GD. 2016. Leaf vein fraction influences the Peclet effect and ^{18}O enrichment in leaf water. *Plant, Cell & Environment* 39: 2414–2427.
- Holloway-Phillips M-M, Cernusak LA, Stuart-Williams H, Ubierna N, Farquhar GD. 2019. Two-source $\delta^{18}\text{O}$ method to validate the CO^{18}O -photosynthetic discrimination model: implications for g_{m} . *Plant Physiology* 181: 1175–1190.
- Horita J, Wesolowski DJ. 1994. Liquid-vapor fractionation of oxygen and hydrogen isotopes of water from the freezing to the critical temperature. *Geochimica et Cosmochimica Acta* 58: 3425–3437.
- Kahmen A, Hoffmann B, Schefuss E, Arndt SK, Cernusak LA, West JB, Sachse D. 2013a. Leaf water deuterium enrichment shapes leaf wax n -alkane dD values of angiosperm plants II: observational evidence and global implications. *Geochimica et Cosmochimica Acta* 111: 50–63.
- Kahmen A, Sachse D, Arndt SK, Tu KP, Farrington H, Vitousek PM, Dawson TE. 2011. Cellulose $\delta^{18}\text{O}$ is an index of leaf-to-air vapor pressure difference (VPD) in tropical plants. *Proceedings of the National Academy of Sciences, USA* 108: 1981–1986.
- Kahmen A, Schefuss E, Sachse D. 2013b. Leaf water deuterium enrichment shapes leaf wax n -alkane dD values of angiosperm plants I: experimental evidence and mechanistic insights. *Geochimica et Cosmochimica Acta* 111: 39–49.
- Kim K, Lee X. 2011. Transition of stable isotope ratios of leaf water under simulated dew formation. *Plant, Cell & Environment* 34: 1790–1801.
- Ladd SN, Maloney AE, Nelson DB, Prebble M, Camperio G, Sear DA, Hassall JD, Langdon PG, Sachs JP, Dubois N. 2021. Leaf wax hydrogen isotopes as a hydroclimate proxy in the tropical Pacific. *Journal of Geophysical Research Biogeosciences* 126: e2020JG005891.
- Lai C-T, Ehleringer JR. 2011. Deuterium excess reveals diurnal sources of water vapor in forest air. *Oecologia* 165: 213–223.
- Lai C-T, Ometto JPHB, Berry JA, Martinelli LA, Domingues TF, Ehleringer JR. 2008. Life form-specific variations in leaf water oxygen-18 enrichment in Amazonian vegetation. *Oecologia* 157: 197–210.
- Lehmann MM, Goldsmith GR, Mirande-Ney C, Weigt RB, Schönbeck L, Kahmen A, Gessler A, Siegwolf RTW, Saurer M. 2020. The ^{18}O -signal transfer from water vapour to leaf water and assimilates varies among plant species and growth forms. *Plant, Cell & Environment* 43: 510–523.
- Lehmann MM, Vitali V, Schuler P, Leuenberger M, Saurer M. 2021. More than climate: hydrogen isotope ratios in tree rings as novel plant physiological indicator for stress conditions. *Dendrochronologia* 65: 125788.
- Liu HT, Schäufele R, Gong XY, Schnyder H. 2017. The $\delta^{18}\text{O}$ and $\delta^2\text{H}$ of water in the leaf growth-and-differentiation zone of grasses is close to source water in both humid and dry atmospheres. *New Phytologist* 214: 1423–1431.
- Liu W, Yang H. 2008. Multiple controls for the variability of hydrogen isotopic compositions in higher plant n -alkanes from modern ecosystems. *Global Change Biology* 14: 2166–2177.
- Mamonov AB, Coalson RD, Zeidel ML, Mathai JC. 2007. Water and deuterium oxide permeability through aquaporin 1: mD predictions and experimental verification. *Journal of General Physiology* 130: 111–116.
- Merlivat L. 1978. Molecular diffusivities of H_2^{16}O , HD^{16}O , and H_2^{18}O in gases. *Journal of Chemical Physics* 69: 2864–2871.
- Munksgaard NC, Cheesman AW, English NB, Zwart C, Kahmen A, Cernusak LA. 2017. Identifying drivers of leaf water and cellulose stable isotope enrichment in *Eucalyptus* in northern Australia. *Oecologia* 183: 31–43.
- Munksgaard NC, Zwart C, Haig J, Cernusak LA, Bird MI. 2020. Coupled rainfall and water vapour stable isotope time series reveal tropical atmospheric processes on multiple timescales. *Hydrological Processes* 34: 111–124.
- Ogée J, Cuntz M, Peylin P, Bariac T. 2007. Non-steady-state, non-uniform transpiration rate and leaf anatomy effects on the progressive stable isotope enrichment of leaf water along monocot leaves. *Plant, Cell & Environment* 30: 367–387.
- Roden JS, Ehleringer JR. 1999. Observations of hydrogen and oxygen isotopes in leaf water confirm the Craig-Gordon model under wide-ranging environmental conditions. *Plant Physiology* 120: 1165–1173.
- Roden JS, Ehleringer JR. 2000a. Hydrogen and oxygen isotope ratios of tree ring cellulose for field-grown riparian trees. *Oecologia* 123: 481–489.
- Roden JS, Ehleringer JR. 2000b. There is no temperature dependence of net biochemical fractionation of hydrogen and oxygen isotopes in tree-ring cellulose. *Isotopes in Environmental and Health Studies* 36: 303–317.
- Roden JS, Lin GG, Ehleringer JR. 2000. A mechanistic model for interpretation of hydrogen and oxygen isotope ratios in tree-ring cellulose. *Geochimica et Cosmochimica Acta* 64: 21–35.
- Rozanski K, Araguas-Araguas L, Gonfiantini R. 1993. Isotopic patterns in modern global precipitation. In: Swart PK, Lohmann KC, McKenzie J, Savin S, eds. *Climate change in continental isotopic records*. Washington, DC, USA: American Geophysical Union, 1–36.
- Sachse D, Billault I, Bowen GJ, Chikaraishi Y, Dawson TE, Feakins SJ, Freeman KH, Magill CR, McInerney FA, van der Meer MTJ *et al.* 2012. Molecular paleohydrology: interpreting the hydrogen- isotopic composition of lipid biomarkers from photosynthesizing organisms. *Annual Review of Earth and Planetary Sciences* 40: 221–249.
- Seibt U, Wingate L, Berry JA, Lloyd J. 2006. Non-steady state effects in diurnal ^{18}O discrimination by *Picea sitchensis* branches in the field. *Plant, Cell & Environment* 29: 928–939.
- Twining J, Stone D, Tadros C, Henderson-Sellers A, Williams A. 2006. Moisture Isotopes in the biosphere and atmosphere (MIBA) in Australia: a priori estimates and preliminary observations of stable water isotopes in soil, plant and vapour for the Tumbarumba Field Campaign. *Global and Planetary Change* 51: 59–72.
- Uemura R, Matsui Y, Yoshimura K, Motoyama H, Yoshida N. 2008. Evidence of deuterium excess in water vapor as an indicator of ocean surface conditions. *Journal of Geophysical Research: Atmospheres* 113: D19114.
- Voelker SL, Brooks JR, Meinzer FC, Roden J, Pazdur A, Pawelczyk S, Hartsough P, Snyder K, Plavcová L, Šantrůček J. 2014. Reconstructing relative humidity from plant $\delta^{18}\text{O}$ and dD as deuterium deviations from the global meteoric water line. *Ecological Applications* 24: 960–975.
- Welp LR, Lee X, Kim K, Griffis TJ, Billmark KA, Baker JM. 2008. $\delta^{18}\text{O}$ of water vapour, evapotranspiration and the sites of leaf water evaporation in a soybean canopy. *Plant, Cell & Environment* 31: 1214–1228.
- Welp LR, Lee X, Griffis TJ, Wen X-F, Xiao W, Li S, Sun X, Hu Z, Val Martin M, Huang J. 2012. A meta-analysis of water vapor deuterium-excess in the midlatitude atmospheric surface layer. *Global Biogeochemical Cycles* 26: GB3021.
- West JB, Sobek A, Ehleringer JR. 2008. A simplified GIS approach to modeling global leaf water isoscapes. *PLoS ONE* 3: e2447.

- Yakir D. 1992. Variations in the natural abundance of oxygen-18 and deuterium in plant carbohydrates. *Plant, Cell & Environment* **15**: 1005–1020.
- Yakir D. 1998. Oxygen-18 of leaf water: a crossroad for plant associated isotopic signals. In: Griffiths H, ed. *Stable isotopes: integration of biological, ecological, and geochemical processes*. Oxford, UK: BIOS Scientific, 147–188.
- Yakir D, Deniro MJ, Rundel PW. 1989. Isotopic inhomogeneity of leaf water: evidence and implications for the use of isotopic signals transduced by plants. *Geochimica et Cosmochimica Acta* **53**: 2769–2773.
- Zech M, Tuthorn M, Detsch F, Rozanski K, Zech R, Zöller L, Zech W, Glaser B. 2013. A 220ka terrestrial $\delta^{18}\text{O}$ and deuterium excess biomarker record from an eolian permafrost paleosol sequence, NE-Siberia. *Chemical Geology* **360–361**: 220–230.
- Zhao L, Wang L, Liu X, Xiao H, Ruan Y, Zhou M. 2014. The patterns and implications of diurnal variations in the d-excess of plant water, shallow soil water and air moisture. *Hydrology and Earth System Sciences* **18**: 4129–4151.
- Zhao LJ, Wang LX, Cernusak LA, Liu XH, Xiao HL, Zhou MX, Zhang SQ. 2016. Significant difference in hydrogen isotope composition between xylem and tissue water in *Populus euphratica*. *Plant, Cell & Environment* **39**: 1848–1857.
- Zundel G, Miekeley W, Grisi BM, Förstel H. 1978. The H_2^{18}O enrichment in the leaf water of tropic trees: comparison of species from the tropical rain forest and the semi-arid region in Brazil. *Radiation and Environmental Biophysics* **15**: 203–212.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

Dataset S1 An EXCEL file containing the compiled dataset that was analysed for this paper.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.