



Michigan Technological University  
*Create the Future* Digital Commons @ Michigan Tech

---

Dissertations, Master's Theses and Master's  
Reports - Open

Dissertations, Master's Theses and Master's  
Reports

---

2013

## Plant-mediated effects on microbial diversity in mesocosms of an oligotrophic bog

Karl J. Romanowicz IV  
*Michigan Technological University*

Follow this and additional works at: <https://digitalcommons.mtu.edu/etds>

 Part of the [Ecology and Evolutionary Biology Commons](#), and the [Microbiology Commons](#)


Copyright 2013 Karl J. Romanowicz IV

---

### Recommended Citation

Romanowicz, Karl J. IV, "Plant-mediated effects on microbial diversity in mesocosms of an oligotrophic bog", Master's Thesis, Michigan Technological University, 2013.  
<https://doi.org/10.37099/mtu.dc.etds/626>

Follow this and additional works at: <https://digitalcommons.mtu.edu/etds>

 Part of the [Ecology and Evolutionary Biology Commons](#), and the [Microbiology Commons](#)

PLANT-MEDIATED EFFECTS ON MICROBIAL DIVERSITY IN MESOCOSMS  
OF AN OLIGOTROPHIC BOG

By  
Karl J. Romanowicz IV

A THESIS  
Submitted in partial fulfillment of the requirements for the degree of  
MASTER OF SCIENCE  
In Applied Ecology

MICHIGAN TECHNOLOGICAL UNIVERSITY

2013

Copyright © Karl J. Romanowicz 2013

This thesis has been approved in partial fulfillment of the requirements for the Degree of MASTER OF SCIENCE in Applied Ecology.

School of Forest Resources and Environmental Science

Thesis Advisor: *Erik A. Lilleskov*

Committee Member: *Rodney A. Chimner*

Committee Member: *Sarah A. Green*

Committee Member: *Jay T. Lennon*

School Dean: *Terry L. Sharik*

## TABLE OF CONTENTS

List of Tables .....	iv
List of Figures .....	v
List of Supplemental Tables .....	vi
List of Supplemental Figures .....	vi
 Introductory Chapter .....	 1
 Chapter 1. Plant-Mediated Effects on Microbial Diversity in Mesocosms of an Oligotrophic Bog .....	  4
1.1 Abstract .....	4
1.2 Introduction.....	5
1.3 Methods and Materials.....	9
1.4 Results.....	15
1.5 Discussion.....	22
1.6 Acknowledgements.....	27
1.7 Literature Cited .....	27
1.8 Tables .....	37
1.9 Figures .....	42
1.10 Supplemental.....	55

## LIST OF TABLES

<b>Table 1</b> Oxidative and hydrolytic extracellular enzymes assayed in bulk peat at the 10-20 cm depth with corresponding enzyme commission number (EC), role in nutrient cycling, and substrate. ....	37
<b>Table 2</b> Bacterial and archaeal diversity indicated by Chao1 richness and equitability evenness in plant functional group treatments. ....	38
<b>Table 3</b> Mean ( $\pm$ SE) percent relative abundance of bacterial phyla and archaeal orders between plant functional groups within sampling depths. ....	39
<b>Table 4</b> Peat soil biogeochemistry, relative percentage of vegetation, and extracellular enzyme activities by plant functional group treatments within depths..	40
<b>Table 5</b> Statistical results for envfit function utilizing environmental data for bacteria and archaea at both 10-20 cm and 30-40 cm depths. ....	41

## LIST OF FIGURES

<b>Figure 1</b> Rank abundance curves for relative abundance of A) bacterial and B) archaeal OTUs obtained from Illumina sequencing .....	42
<b>Figure 2</b> Water table depth in mesocosm bins for high water table treatment (HWT) and low water table treatment (LWT) during the two-week peat coring period. ....	43
<b>Figure 3</b> Bacteria phylum-level community composition between sampling depth..	44
<b>Figure 4</b> Bacteria phylum-level community composition between plant functional group treatments within sampling depths .....	45
<b>Figure 5</b> Archaea order-level community composition between sampling depth. ....	46
<b>Figure 6</b> Archaea order-level community composition between plant functional group treatments within sampling depths. ....	47
<b>Figure 7</b> Heatmap visualization for top 100 most abundant bacterial OTUs indicating relative abundance by depth.....	48
<b>Figure 8</b> Heatmap visualization for all archaeal OTUs indicating relative abundance by depth.....	49
<b>Figure 9</b> Principal coordinates analysis of A) bacteria and B) archaea. ....	50
<b>Figure 10</b> PCoA of top 100 OTUs with envfit environmental vectors for A) bacteria at 10-20 cm depth and B) bacteria at 30-40 cm depth. ....	51
<b>Figure 11</b> PCoA of all OTUs with envfit environmental vectors for A) archaea at 10-20 cm depth and B) archaea at 30-40 cm depth.....	52
<b>Figure 12</b> Mean relative abundance of all methanotrophic bacteria summed between sampling depths .....	53
<b>Figure 13</b> Mean relative abundance of all methanogenic archaea summed between sampling depths .....	54

## LIST OF SUPPLEMENTAL TABLES

<b>Table S1</b> Depth effects on top 50 most abundant bacterial OTUs as calculated through the OTU Category Significance ANOVA in QIIME. ....	55
-----------------------------------------------------------------------------------------------------------------------------------------------	----

<b>Table S2</b> Depth effects on archaeal OTUs as calculated through the OTU Category Significance ANOVA in QIIME and list of all archaeal OTUs with consensus lineage from the oligotrophic bog mesocosm study. ....	56
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----

## LIST OF SUPPLEMENTAL FIGURES

<b>Figure S1</b> Electrophoresis DNA quantity visualization on 2:1% agarose gel for DNA extractions at the 10-20 cm depth. ....	57
---------------------------------------------------------------------------------------------------------------------------------	----

<b>Figure S2</b> Electrophoresis DNA quantity visualization on 2:1% agarose gel for DNA extractions at the 30-40 cm depth. ....	58
---------------------------------------------------------------------------------------------------------------------------------	----

<b>Figure S3</b> Relative abundance of all known methanotrophic bacteria within plant functional group treatments between sampling depths as found in this study. ....	59
------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----

<b>Figure S4</b> Mean relative abundance of all known methanogenic archaea plus unclassified Euryarchaeota within plant functional group treatments between sampling depths as found in this study. ....	60
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----

## INTRODUCTORY CHAPTER

Northern peatlands form in moderately cool yet wet climatic zones where photosynthetic capacity of vegetation exceeds decomposition. Accumulating slowly throughout the Holocene, these peat deposits now cover approximately 3% of terrestrial land area and store up to one-third of global terrestrial carbon stocks. This vast carbon store has the potential to contribute substantial feedbacks to global climate change, as the balance between carbon inputs from vegetation and carbon loss from decomposition is relatively sensitive to changing climate. Changes in plant communities may lead to feedbacks altering the carbon balance of peatlands. No study to date has tested the direct effects of dominant plant functional group manipulations within an oligotrophic bog on rhizosphere microbial communities and their associated functions.

The two major vascular plant functional groups in northern peatlands are sedges and shrubs of the family Ericaceae. Specific traits of these two functional groups are hypothesized to drive opposing effects on peatland carbon balance. Sedge species contain aerenchyma that function as a ventilation system by transporting oxygen to the root tips in saturated, anaerobic soil while simultaneously transporting carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>) from the soil to the atmosphere. As a result of aerenchyma, sedges serve as direct channels that bypass oxidizing environments and influence rhizospheric processes regardless of water table levels. Sedges are hypothesized to create extensive carbon oxidation and mineralization hotspots throughout the peat profile and may lead to enhanced aerobic microbial activity near root surfaces promoting increased peat subsidence and negative carbon balance. In contrast, ericaceous shrubs are hypothesized to drive decreased carbon mineralization by mediating changes in rhizosphere microbial communities and enzyme activity. Ericaceous shrubs lack aerenchyma, have poor litter quality, and utilize enzymatically active ericoid mycorrhizal fungi that may suppress free-living saprotrophs through



direct interception of necessary macronutrients such as nitrogen and phosphorous. This potential combination of decreased carbon mineralization and increased net primary production (NPP) could drive continued positive carbon balance.

This thesis documents the response of rhizosphere microbial communities to changes in plant functional groups in a mesocosm-based experimental design. Research was conducted at the Houghton Mesocosm Facility at the USDA Forest Service Northern Research Station Forestry Sciences Laboratory in Houghton, Mich., USA. Peat monoliths were harvested and housed in mesocosm chambers in May 2010 and underwent plant functional groups manipulations in July 2011. Mesocosms were manipulated into three vegetation treatments: unmanipulated (+sedge, +Ericaceae), sedge (+sedge, -Ericaceae), and Ericaceae (-sedge, +Ericaceae). The Houghton Mesocosm Facility provided fine-scale control of plant community composition allowing us to investigate how carbon-based rhizospheric processes in peatland ecosystems would respond under predicted future climate change scenarios.

Our research explored the impacts of plant functional groups on rhizosphere microbial community composition at two depths from cores collected during September 2011. These samples represent microbial community composition approximately two months following plant functional group manipulations under anaerobic soil conditions created by high water table levels. Microbial DNA was extracted from both 10-20 cm and 30-40 cm depths (both depths were below water table levels at time of sampling) from each mesocosm core and sequenced for bacterial and archaeal community composition through 16S rRNA targeted gene sequencing on an Illumina MiSeq platform.

Results indicated significant differences in bacterial and archaeal community composition between sampling depths. Heterogeneity within bacterial community

composition decreased with increasing depth in the peat soil, and was paralleled by an increase in the relative abundance of anaerobic taxa. Plant functional groups had no significant influence at the 30-40 cm depth, however, within the more shallow 10-20 cm depth, plant functional groups significantly influenced certain bacterial and archaeal communities. These results implied increased microbial heterogeneity within the 10-20 cm depth under the Ericaceae treatment, incorporating a more even distribution of Acidobacteria, Bacteroidetes and Gammaproteobacteria. Sedge treatments experienced an increase in the relative dominance of Acidobacteria, while also encouraging significant increases in Alphaproteobacteria, and Verrucomicrobia, which are known to contain methane-oxidizing taxa and warranted a more thorough examination. Sampling depth significantly influenced the relative abundance of both methane-oxidizing bacteria and methanogenic archaea, but there were no significant effects of plant functional groups. These results suggest that the bacteria and archaea involved in methane oxidation or production, respectively, are not directly influenced by plant functional groups following such a short timeframe from manipulations.

Although these are early results in a long-term mesocosm study, they provide essential information regarding initial shifts in the relative abundance of microbial community composition under specific plant functional group manipulations. Results from our mesocosm-based study have begun to test whether plant functional groups have the potential to significantly alter the mechanisms controlling carbon-based rhizospheric processes by fundamentally changing microbial community composition.

# **CHAPTER 1. PLANT-MEDIATED EFFECTS ON MICROBIAL DIVERSITY IN MESOCOSMS OF AN OLIGOTROPHIC BOG**

## **1.1 Abstract**

Globally, peatlands occupy a small portion of terrestrial land area but contain up to one-third of all soil organic carbon. This carbon pool is vulnerable to increased decomposition under projected climate change scenarios but little is known about how plant functional groups will influence microbial communities responsible for regulating carbon cycling processes. Here we examined initial shifts in microbial community structure within two sampling depths under plant functional group manipulations in mesocosms of an oligotrophic bog. Microbial community composition for bacteria and archaea was characterized using targeted 16S rRNA Illumina gene sequencing. We found statistically distinct spatial patterns between the more shallow 10-20 cm sampling depth and the deeper 30-40 cm depth. Significant effects by plant functional groups were found only within the 10-20 cm depth, indicating plant-mediated microbial community shifts respond more quickly near the peat surface. Specifically, the relative abundance of Acidobacteria decreased under ericaceous shrub treatments in the 10-20 cm depth and was replaced by increased abundance of Gammaproteobacteria and Bacteroidetes. In contrast, the sedge rhizosphere continued to be dominated by Acidobacteria but also promoted an increase in the relative recovery of Alphaproteobacteria and Verrucomicrobia. These initial results suggest microbial communities under ericaceous shrubs may be limited by anaerobic soil conditions accompanying high water table conditions, while sedge aerenchyma may be promoting aerobic taxa in the upper peat rhizosphere regardless of ambient soil oxygen limitations.

## 1.2 Introduction

Northern peatlands store approximately a third of all terrestrial soil carbon in the form of partially decayed organic peat (Gorham, 1991; Clymo et al., 1998) while also emitting 9-18% of all global atmospheric methane (Laanbroek, 2009). Carbon storage is a result of high water table levels and subsequent oxygen limitation in peat soils that favors primary production over decomposition (Clymo, 1984). The unique hydrology of peatlands promotes anaerobic soil conditions that stimulate methanogenic microorganisms to produce methane ( $\text{CH}_4$ ) as the terminal step in anoxic microbial degradation of organic matter (Sundh et al., 1994). Historically, peatlands have served as net carbon sinks but could transition to carbon sources if water tables become lowered as a result of climate change or anthropogenic drainage (Trettin et al., 2006; Dinsmore et al., 2010). Shifts in the environmental and physico-chemical constraints on microbial decomposition could alter carbon cycling and methane production/consumption processes (Andersen et al., 2012). This could lead to increased peat oxidation to carbon dioxide ( $\text{CO}_2$ ), while limiting methane production as anaerobic soil conditions decline. However, plant functional groups have the ability to regulate soil microbial processes by influencing litter quality (Joannis et al., 2007), oxygen availability (Strack et al., 2006), root exudation (Inderjit & Mallik, 2002), and nutrient competition (Bonfante & Genre, 2010; Artz et al., 2007). To our knowledge, no study has been conducted to specifically test the direct consequences of spatial and temporal changes to peatland plant functional groups on microbial community structure and function.

Bogs are oligotrophic peatlands deriving limited nutrient inputs mainly from precipitation (Crum, 1991). Northern regions have experienced increasing variability in precipitation patterns in recent decades (Tsonis et al., 1996; Groisman et al., 2005) causing declines in peatland water tables and increased frequency and intensity of mid-summer water table drawdown (Roulet et al., 1992; Hilbert et al., 2000).

Sustained changes in peatland water table levels have been shown to directly influence the establishment of plant functional groups (Breeuwer et al., 2009; Weltzin et al., 2000). Dominant groups of bog plant communities include *Sphagnum* mosses, sedges, and ericaceous shrubs (Crum, 1991). *Sphagnum* lack root systems and form dense lawns on the surface of peatlands (Strack et al., 2006) where they attract water to the surface through capillary forces (Farrick, 2009). Sedges are perennial graminoids that use aerenchyma to promote diffusion of oxygen to deep roots in anoxic peat and subsequently create extensive carbon oxidation and mineralization hotspots (Holzapfel-Pschorn et al., 1986). In contrast, ericaceous shrubs are shallowly rooted and form a symbiotic relationship with enzymatically active ericoid mycorrhizal fungi (Cairney & Burke, 1998) possibly suppressing free-living heterotroph activity and promoting decreased carbon mineralization. In essence, root traits are predicted to be dominant regulators of microbially mediated soil processes as they alter carbon and nutrient resources and influence rhizosphere conditions suitable for specific functional groups of microorganisms (Andersen et al., 2012; Strakova et al., 2011).

Acidic, nutrient-poor conditions in bogs are a major constraint on bacterial community composition (Dedysh et al., 1998), along with energetic constraints such as decreased availability of oxygen and high quality electron acceptors with increasing depth in the peat column (Artz, 2009). However, plant functional groups have the ability to modify environmental conditions and subsequently alter microbial composition such that sedge dominated peatlands can harbor significantly different communities from ericaceous shrub peatlands (Fisk et al., 2003; Borga et al., 1994). Archaeal community structure also seems to be substrate and plant specific as observed differences were found in community composition among plant species within the rhizosphere (Cadillo-Quiroz et al., 2009). Ericoid mycorrhizal fungi are a prime example of a plant-dependent microbe and are hypothesized to dominate upper surface layers in ericaceous shrub peatlands, promoting enzyme activity primarily

focused on accessing non-carbon nutrients such as organically bound nitrogen and phosphorus for the fungal and plant partners (Cairney & Burke, 1998). As a result, the saprotrophic activity of ericoid mycorrhizal fungi could permit them to be effective competitors for nutrients with free-living heterotrophs, suppressing their activity to avoid microbial consumption of depolymerized substrates (Allison, 2005) and potentially limiting carbon mineralization activity (Lindahl et al., 2002). In contrast, sedge rhizospheres are substantially different environments due to an absence of mycorrhizas, deep root penetration into the peat, and abundance of oxygen supplied via aerenchyma, which may support aerobic microorganisms near root surfaces throughout the peat rhizosphere.

Plant functional group influences on peatland microbial community structure and function have previously been studied through field-based and mesocosm experiments contrasting oligotrophic bogs with minerotrophic fen peatlands (*see* Lin et al., 2012; Strakova et al., 2011; Myers et al., 2011; Peltoniemi et al., 2009; Kim et al., 2008; Jaatinen et al., 2007; Sowerby et al., 2005). However, bogs and fens fundamentally differ not only in plant functional group composition but also through hydrology, pH, nutrient availability, and soil chemistry (Bridgham et al., 1996) to such an extent that direct effects of plant functional groups are difficult to distinguish from myriad other environmental and physico-chemical conditions. In this study, we developed an experimental mesocosm facility to control for these environmental variables. We utilized oligotrophic bog monoliths originally composed of relatively equal distributions of sedge and ericaceous shrubs, and implemented vascular plant functional group manipulations to create either sedge dominant, ericaceous shrub dominant, or mixed dominance systems. To test our hypotheses, bacterial and archaeal communities were sequenced to determine changes in community composition following plant functional group manipulations. As microbial communities within peat monoliths were originally identical under the mixed plant

system, any shift in community composition can be considered a direct result of the plant functional group manipulations.

The objective of this study was to quantitatively evaluate the initial shifts in microbial community structure in response to plant functional group manipulations. We hypothesized that the dominant vascular plant communities, sedges and ericaceous shrubs, would have different effects on peatland microbial community composition. We expected under high water table conditions an increase in the relative abundance of aerobic organic matter decomposers (e.g. taxa within Actinobacteria and Bacteroidetes phyla) and methanotrophic (methane oxidizing) bacteria (e.g. taxa within Alpha- and Gammaproteobacteria and Verrucomicrobia) throughout the sedge rhizosphere due to increased oxygen availability via aerenchyma. In contrast, the Ericaceae rhizosphere should experience an increase in the relative abundance of Acidobacteria (low C mineralization rates) and anaerobic-tolerant taxa as well as an increase in the relative abundance of methanogenic (methane producing) archaea at the deeper anoxic depths. Regardless of plant functional groups, we hypothesize a greater relative abundance of aerobic microorganisms at the more shallow 10-20 cm depth with increasing abundance of anaerobic taxa and archaea especially at the deeper 30-40 cm depth due to limitations in soil oxygen availability with increasing depth. Such changes in microbial community structure due to plant functional group manipulations are hypothesized to promote peat oxidation, carbon loss, and decreased methane production from sedge-dominated peatlands while Ericaceae peatlands would limit the effects of peat oxidation and organic carbon decomposition and support natural diffusion of methane to the atmosphere while limiting methane oxidation in the rhizosphere.

### 1.3 Methods and Materials

#### 1.3.1 Site Description

Peat monoliths were collected in close proximity to each other in May 2010 from an acidic (pH 4.0), oligotrophic bog in northeastern Minnesota (47°07'05"N, 92°47'59"W, Meadowlands, MN). Vegetation included bryophytes (*Sphagnum fuscum* (Schimp.) Klinggr., *S. magellanicum* Brid., *S. rubellum* Wilson, *Polytrichum strictum* Brid.), sedges (*Carex oligosperma* Michx., *Eriophorum angustifolium* Honck.), and ericaceous shrubs (*Andromeda glaucophylla* Link., *Chamaedaphne calyculata* (L.) Moench., *Kalmia polifolia* Wang., *Rhododendron groenlandicum* Oeder., and *Vaccinium oxycoccos* L.). Peat monoliths were extracted from the treeless portion of the bog with *a priori* designation of relatively equal representation of both sedge and Ericaceae vascular plant cover (*results not shown*), while *E. angustifolium* was specifically avoided due to sparse and variable distribution.

#### 1.3.2 Experimental Design

Twenty-four 1-m<sup>3</sup> peat monoliths were extracted from the bog, housed in Teflon-coated stainless steel mesocosm chambers, transported to Houghton Mich., and placed in the Houghton Mesocosm Facility at the USDA Forest Service Forestry Sciences Laboratory. Each mesocosm has been implemented with two 80 cm long temperature probes with 5 thermistors each to monitor vertical and horizontal temperature gradients, a dissolved oxygen sensor, a pressure transducer to continuously measure water table depth, with the latter linked to a proportional control valve attached to a porewater intake manifold to regulate and collect outflow water when water tables are sufficiently high to initiate flow. Beginning in July 2011, individual mesocosm chambers were randomly assigned one of the following plant functional group treatments: *i.*) “Unmanipulated” – sedge and Ericaceae left intact;



ii.) “Sedge” – sedge left intact while Ericaceae removed; and iii.) “Ericaceae” – Ericaceae left intact while sedge removed. Plant functional group treatments were replicated eight times and randomly assigned to establish the experimental design (3 plant functional group x 8 replicates,  $N = 24$ ).

To manipulate plant functional groups in July 2011, all above- and easily removable belowground biomass (buried stems) of Ericaceae was carefully removed while minimizing disturbance. To avoid significant peat disturbance only aboveground biomass of sedges was removed. Subsequent removals of regrown targeted biomass were implemented by clipping weekly throughout the growing season to ensure complete removal of the targeted plant communities. Along with plant functional group treatments, each mesocosm chamber was assigned either a “high” or “low” water table treatment with seasonal profiles based on typical low variability, high minimum water table years (high treatment), and typical high variability, low water table years (low treatment) derived from the almost 50 year record of precipitation and water tables at the Marcell Experimental Forest (Kolka et al., 2011). In year 1 of the vegetation treatments (2011, year of present study) water table manipulations were kept to a minimum (approximately 5 cm difference between high and low water treatments during summer) to permit full vegetation recovery and were not considered as a treatment combination (e.g. Sedge – High water table) in the experimental design until 2012.

### *1.3.3 Peat Sampling, Biogeochemical Properties, and Enzyme Assays*

An individual peat core (0-80 cm depth; 3 cm diameter) was collected from each mesocosm chamber ( $N=24$ ) over a two-week sampling period spanning August 31 to September 13, 2011. These cores represented the vertical peat soil profile approximately two months following plant functional group manipulations. Each core was partitioned into 10 cm vertical increments, placed in labeled Ziploc bags,

and immediately stored at -80°C to preserve microbial nucleic acids for future analysis. The 10-20 cm and 30-40 cm vertical increments were utilized for all subsequent microbial analyses.

Porewater samples were collected at the same depths as the peat samples (20 cm and 40 cm) by Aleta Daniels (graduate student – Michigan Tech) using piezometers. Laboratory techniques for analyzing dissolved organic carbon (DOC) and total dissolved nitrogen (TDN) concentrations were described previously using high temperature catalytic oxidation techniques (Olefelt & Roulet, 2012) with modifications specific for this study introduced by Evan Kane (personal communication). In addition, two optical properties indicating alterations in DOC composition were analyzed. The first property, specific ultraviolet absorbance (SUVA<sub>254</sub>), was calculated by normalizing UV absorbance at 254 nm against DOC concentrations. The SUVA<sub>254</sub> index is reported to increase linearly with measured DOC aromaticity (Weishaar et al., 2003). The second property, E<sub>2</sub>:E<sub>3</sub>, is the UV absorbance ratio between 254 nm and 365 nm. The E<sub>2</sub>:E<sub>3</sub> ratio decreases as molecular size increases allowing for an estimation of dissolved organic matter (DOM) size (De Haan & De Boer, 1987).

Extracellular enzyme assays were conducted using methods described by Saiya-Cork et al. (2002), with modifications for organic soil. All enzyme assays were conducted within 4 hours of peat core collection on a vertical split of the 10-20 cm core increment. Enzyme assays were not conducted on the 30-40 cm depth samples. In brief, bulk peat samples from the 10-20 cm depth were prepared in suspensions by adding 1.0 g (wet weight) peat to 125 ml of deionized water and shaken for two minutes to homogenize thoroughly. Peat suspensions were stirred continuously on a magnetic stir plate while 200 µL aliquots were dispensed into 96-well microplates with eight replicate wells per sample per assay. The peat suspensions were assayed for two oxidative enzyme activities and three hydrolytic enzyme activities involved

with carbon, sulfur, and phosphorous cycling (Table 1). All hydrolytic assays were analyzed fluorimetrically with 365 nm excitation and 450 nm emission filters, while oxidative assays were measured spectrophotometrically by quantifying absorbance at 450 nm. Both fluorimetric and absorbance measurements were performed on a SpectraMax M2 plate reader (Molecular Devices, Sunnyvale, California). All subsequent plating conditions were followed according to Saiya-Cork et al. (2002) with the following modifications: eight replicate wells for each blank, negative control, and quench standard, and all microplates were incubated in the dark at room temperature for 3 hours or 24 hours for hydrolytic or oxidative assays, respectively. Also, 50  $\mu$ L of 5 mM L-3,4-dihydroxyphenylalanine (L-DOPA) were added to each sample well for measuring phenol oxidase and peroxidase enzyme activities as modified from 25 mM L-DOPA for ease of dissolving substrate in deionized water. Comparison of measured enzyme activity between 5 mM and 25 mM L-DOPA showed no statistical differences (*results not shown*). Potential oxidative enzyme activity for peat suspensions were expressed as 1  $\mu$ mol of L-DOPA substrate oxidized  $\text{h}^{-1} \text{g}^{-1}$  dry mass. Potential hydrolytic enzyme activity for peat suspensions were expressed as 1 nmol of substrate consumed  $\text{h}^{-1} \text{g}^{-1}$  dry mass.

#### *1.3.4 DNA Extraction and Illumina Sequencing*

Microbial DNA extractions were carried out at the Forestry Sciences Laboratory in Houghton, Michigan at two depths from peat cores collected in September 2011. In brief, all peat core increments from both the 10-20 cm depth and 30-40 cm depth were removed from -80°C storage, suspended in liquid nitrogen, fragmented by hand in mortar and pestle, re-suspended in liquid nitrogen, then finely ground in an electric coffee grinder prior to DNA extractions (sterilized between samples). Following homogenization, total genomic DNA was extracted from 0.5 g peat soil using a MoBio PowerSoil® DNA Isolation kit following the manufacturer's instructions (MoBio Laboratories Inc., Carlsbad, California). Extracted DNA quantity and quality

visualizations were performed on 2:1% agarose gel through electrophoresis for 10-20 cm depth (Supplemental Figure S1) and 30-40 cm depth (Supplemental Figure S2). Isolated DNA samples were purified using a MoBio PowerClean® DNA Clean-Up kit following manufacturer's instructions (MoBio Laboratories Inc., Carlsbad, California) and stored in TE buffer solution pH 7.6. DNA quantitation was assessed on a NanoDrop 1000 spectrophotometer (Thermo Scientific Inc., Waltham, Massachusetts) to ensure sample quantities exceeded 10 ng  $\mu\text{L}^{-1}$  (*results not shown*). All purified DNA samples were sent to U.S. Department of Energy Joint Genome Institute (JGI) in Walnut Creek, California for amplification and iTAG 16S V4 region sequencing (*see* Caporaso et al., 2011) on the Illumina MiSeq platform.

### *1.3.5 Illumina Sequencing Analysis Pipeline*

All 16S rDNA amplicons (iTags) were sequenced on the Illumina MiSeq instrument with a 2x250 bp reads configuration resulting in roughly 4 million sequenced forward reads and 4 million reverse reads. Julien Tremblay performed the following methods at the Joint Genome Institute and provided all method descriptions. In brief, all unpaired reads following sequencing were discarded and all remaining reads were trimmed to 165 bp. Reads were assembled (overlapping paired assembly) with FLASH software (Magoc & Salzberg, 2011) and primer sequences were removed from the assembled reads. Assembled reads were trimmed from both 5' and 3' ends using a 20 bp sliding window quality threshold having a mean quality of 30. The trimmed assembled reads were filtered for quality where all reads having more than 10 nucleotides below quality 15 were discarded. Remaining reads were referred to as filtered reads and were clustered into operational taxonomic units (OTUs) at 97% similarity with JGI's in-house clustering algorithm (SeqObs). No denoising or chimera removal step was implemented, but singleton OTUs were removed prior to downstream analyses. Remaining OTUs were analyzed for taxonomic distribution using QIIME version 1.6.0 (Caporaso et al., 2010) where OTUs were classified with

the Ribosomal Database Project (RDP) classifier using a JGI in-house training set containing the complete Greengenes database supplemented with eukaryotic sequences from the Silva databases and a customized set of mitochondria and chloroplasts 16S sequences.

### *1.3.6 Statistical Analyses*

Bacterial and archaeal richness and evenness were calculated in QIIME version 1.6.0 based on their original OTU tables rarefied to 6,688 bacterial sequence reads and 23 archaeal sequence reads per sample, respectively. Microbial diversity was expressed as Chao1 richness and equitability evenness (1 = complete equitability). Pairwise sample dissimilarities of microbial community structure were calculated as Bray-Curtis distances with 1,000 permutations, and visualized using principal coordinate analysis (PCoA) ordination. Environmental effects (e.g. porewater chemistry, relative percentage of vegetation, and extracellular enzyme activity) on microbial community variation were evaluated within each depth through the envfit function of the vegan package in R (*see* Oksanen et al., 2011) with 1,000 permutations using the bray-curtis distance matrix derived for PCoA. Plant functional group and soil depth effects on bacterial and archaeal taxonomic diversity at the phylum and order level, respectively, were calculated using multivariate analysis of variance (MANOVA) in R. Post-hoc tests utilizing Tukey's Honestly Significant Difference were conducted on variables considered significant ( $p < 0.05$ ) through MANOVA analysis to determine which treatment combinations created significant differences. OTU category significance tests were performed in QIIME on bacterial and archaeal OTUs utilizing analysis of variance (ANOVA) for categorical variables such as sampling depth and plant functional group, while Pearson correlations were utilized for continuous variables including sedge percent cover, pH, and water table.

## 1.4 Results

### 1.4.1 *Bacterial and Archaeal Sampling Diversity*

The Illumina sequencing bacterial data set contained 1,534,582 total read sequences with an average  $34,101 \pm 15,608$  reads per sample, which clustered into 6,218 OTUs at the 97% sequence similarity level. Likewise, the archaeal data set contained 9,713 total read sequences with an average  $215 \pm 178$  reads per sample, which clustered into 44 OTUs at the 97% sequence similarity level. Bayesian classification indicated 99.37% of total read sequences were bacteria and 0.63% of total read sequences were archaea. All OTU-based analyses were performed on the original OTU table rarefied to the lowest total sample sequence read for both bacteria (6,688 reads) and archaea (23 reads) to minimize the effects of uneven sampling effort in comparing diversity across samples (Table 2).

### 1.4.2 *Bacterial and Archaeal Species Richness and Evenness*

The rank abundance distributions for bacteria showed strong dominance of a few OTUs with a long tail of rare OTUs irrespective of plant functional group treatment or depth. Rank abundance for archaea was similar, indicating ~12 OTUs representing 1% or greater relative abundance followed by ~20-30 OTUs each representing less than 1% of total relative abundance (Figure 1). During the two week peat coring process in September 2011, water table levels remained above the 10-20 cm sampling depth (Figure 2) indicating all samples collected should have been experiencing anaerobic soil conditions. Statistical analyses of bacterial species richness indicated no significant differences between plant functional groups or depth but evenness was significantly different between sampling depths ( $p=0.009$ ) with increased equitability within the 30-40 cm depth (Table 2). Archaeal richness and evenness was not significantly influenced by depth or plant functional group treatments. These results

suggests that greater bacterial species evenness at the deeper 30-40 cm depth indicates greater homogeneity, and confirms that the bacterial community was dominated by a select number of OTUs as alluded to in the rank abundance distribution.

#### *1.4.3 Bacterial and Archaeal Taxonomic Classification*

Bacterial OTUs were assigned to 15 phyla (with Proteobacteria separated into 5 classes) while archaeal OTUs were assigned to 2 phyla with all subsequent analyses comparing 9 orders within 6 classes of the 2 phyla (Table 3). Across all treatment combinations, dominant phyla included (in order of decreasing abundance) Acidobacteria, Bacteroidetes, Alphaproteobacteria, Gammaproteobacteria, Cyanobacteria, Firmicutes, Verrucomicrobia, Actinobacteria, Planctomycetes, and Deltaproteobacteria. The remaining 7 phyla (and 3 Proteobacteria classes) combined represented less than 5% of total community composition and included (in order of decreasing abundance) Chlamydiae, Betaproteobacteria, Proteobacteria unclassified, Spirochaetes, Nitrospirae, Tenericutes, Armatimonadetes, Chloroflexi, Epsilonproteobacteria, and Fusobacteria. 18% of all bacterial OTUs were not classified beyond domain.

Overall, the taxonomic distribution of OTUs within the bog soils indicated certain significant changes in community composition based on both depth (Figure 3) and plant functional group treatments (Figure 4). MANOVA results indicated bacterial community composition significantly differed between the 10-20 cm and 30-40 cm depths and also differed within depths by plant functional group manipulations. Acidobacteria was the dominant phylum throughout the bog soils representing 39% of total community composition and significantly differed between depths ( $p < 0.0001$ ) with greatest relative abundance within the 30-40 cm depth and also differed between sedge and Ericaceae treatments ( $p = 0.020$ ) with greatest relative abundance within

sedge treatments. Likewise, Bacteroidetes, which represented 11% of total bacterial composition, differed between depths ( $p < 0.0001$ ) with greatest relative abundance within the 10-20 cm depth and also differed between sedge and Ericaceae treatments ( $p = 0.012$ ) with greatest relative abundance within the Ericaceae treatments. However, Acidobacteria and Bacteroidetes did not statistically differ between sedge and unmanipulated treatments ( $p = 0.125$ ;  $p = 0.244$ , respectively) or Ericaceae and unmanipulated treatments ( $p = 0.648$ ;  $p = 0.313$ , respectively) suggesting sedge and ericaceous plant communities are driving opposite effects on both Acidobacteria and Bacteroidetes abundance especially at the 10-20 cm depth.

Alphaproteobacteria (9% of total bacteria), Gammaproteobacteria (6% of total bacteria), and Verrucomicrobia (4% of total bacteria) differed only between sampling depths (all  $p < 0.05$ ) but were not significantly influenced by plant functional groups or the interaction between depth and plant functional group treatments. Specifically, the 10-20 cm depth had higher relative abundance of Alpha- and Gammaproteobacteria while the 30-40 cm depth had higher relative abundance of Verrucomicrobia.

A thorough examination of methane oxidizing species indicated that our study included the following taxa: Alphaproteobacteria *Methylocella* spp. of the family Beijerinckiaceae, *Methylocystis* spp. and *Methylosinus* spp. of the family Methylocystaceae; Gammaproteobacteria *Methylococcus* spp. and *Methylomonas* spp. of the family Methylococcaceae; and Verrucomicrobia *Methylacidiphilum* spp. of the family Methylacidiphilaceae (see Supplemental Figure S3). Increasing depth had a significant positive impact on the relative abundance of methanotrophs only in the unmanipulated plant functional group treatment ( $p < 0.05$ ), but trends suggested a greater abundance of these taxa exists within sedge treatments compared to Ericaceae treatments in the 10-20 cm depth (Figure 12).



Dominant archaeal communities included members of the methanogenic Euryarchaeota such as *Methanobacteriales* (23%), *Methanomicrobiales* (11%), *Methanosarcinales* (2%), and *Methanocellales* (<1%) with a large proportion of unclassified Euryarchaeota (22%) (Table 3). Non-methanogenic Crenarchaeota included *Cenarchaeales* (28%), *Thermoproteales* (<1%), and *Nitrososphaerales* (<1%). Over 5% of all archaea sequenced were unclassified beyond domain. *Cenarchaeales* was the dominant non-methanogenic Crenarchaeota across the majority of treatments.

Although total archaeal abundance was low, there were considerable variations between depths (Figure 5) and marginally significant differences between plant functional group treatments (Figure 6). The proportion of methanogenic *Methanobacteriales* at the 10-20 cm depth was more than double the 30-40 cm depth ( $p=0.002$ ) but a large proportion of Euryarchaeota were unclassified at the 30-40 cm depth compared to the 10-20 cm depth ( $p<0.0001$ ) (see Supplemental Figure S4). Likewise, *Methanomicrobiales* was 7 times greater at the 10-20 cm depth compared to the 30-40 cm depth ( $p=0.001$ ). The relative recovery of methanogens overall was influenced by sampling depth with greatest relative abundance within the 10-20 cm depth, while plant functional groups appeared to have no significant influence on relative recovery (Figure 13). It should be noted that the Euryarchaeota member *Halobacteriales* was initially 17 times greater in the Ericaceae treatment at the 10-20 cm depth compared to all other treatment combinations, which was a direct result of an outlier community in one of the replicate mesocosms (i.e. Bin 13, Ericaceae treatment, 10-20 cm depth consisted of 83% *Halobacteriales*) but this sample was removed from all analyses as it was believed to be a contamination issue possibly incorporated during the sequencing process (S. Tringe, personal communication).

#### 1.4.4 Significant Bacterial and Archaeal OTUs

An OTU significance test based on analysis of variance (ANOVA) between sampling depths was performed in QIIME on the top 100 most abundant bacterial OTUs and indicated that 46 of those OTUs had a significant preference for a particular depth after Bonferroni correction (Supplemental Table S1). Within archaea, 6 of 46 OTUs were found to have significant preferences by depth following Bonferroni correction (Supplemental Table S2). Heatmaps displayed visual differences in the relative abundance of the top 100 bacterial OTUs by treatment and depth combination (Figure 7) and all archaeal OTUs by treatment and depth combination (Figure 8).

After determining significant differences between sampling depths, further OTU significance tests were performed within each depth. An OTU significance test based on ANOVA between plant functional groups was performed within both sampling depths for bacteria and archaea, but following Bonferroni corrections, no OTUs were found to have significant plant functional group preferences (*results not shown*). Pearson correlations were calculated utilizing sedge percent cover, pH, and water table on bacterial and archaeal OTUs within both sampling depths but also found no significant results within any treatment combination (*results not shown*).

#### 1.4.5 Community Comparison Across Plant Functional Groups and Depth

Bacterial and archaeal communities separated by depth along the first axis (PCO1) of the principal coordinate analysis based on Bray-Curtis similarity (Figures 9A; 9B). PCO1 for bacteria explained 50.37% of total variation while PCO1 for archaea explained 33.92% of total variation. The second axis (PCO2) of the PCoA separates bacterial and archaeal communities by plant functional groups. PCO2 for bacteria explained 10.94% of total variation while PCO2 for archaea explained 13.69% of

total variation. Depth appears to be the dominant factor driving microbial community variation, however, within the 10-20 cm depth it appears that microbial communities respond differently between sedge and Ericaceae treatments. Taxonomic variation was calculated and plotted alongside plant functional group by depth treatments on the PCoA to further illustrate their effects on altering microbial community composition. Within bacteria, Cyanobacteria and Bacteroidetes appear to have shifted towards the 10-20 cm depth and their relative recovery increased the most within the unmanipulated and Ericaceae treatments. Firmicutes was the only phylum that had a strong preference for the 30-40 cm depth and all remaining phyla had no preference for either depth or any particular plant functional group treatment. For the archaea, methanogenic *Methanobacteriales* and *Methanomicrobiales* were most abundant at the 10-20 cm depth within the unmanipulated and sedge treatments. All subsequent orders except *Cenarchaeales* increased in abundance in the 30-40 cm depth with no noticeable preference under sedge or Ericaceae treatments.

#### 1.4.6 Environmental Influence on Bacterial and Archaeal Community Composition

A combination of data including pH, porewater biogeochemistry, optical properties of DOC, water table levels, relative percentage of vegetation species, and extracellular enzyme activities (Table 4) were incorporated into an envfit analysis against OTU distributions for both bacteria and archaea to better understand correlations between microbial community composition and environmental factors. Results from the envfit analysis (Table 5) for bacteria at the 10-20 cm depth indicated that the relative percentage of *Carex oligosperma* ( $p=0.001$ ) along with DOC concentrations ( $p=0.040$ ), the ratio of  $E_2:E_3$  ( $p=0.034$ ), and sulfatase activity ( $p=0.002$ ) all significantly correlated with community variation (Figure 10A). Likewise, the relative percentage of *Vaccinium oxycoccos* ( $p=0.085$ ), the ratio of  $SUVA_{254}$  ( $p=0.099$ ), and  $\beta$ -glucosidase activity ( $p=0.058$ ) were all marginally significantly correlated with community variation at the 10-20 cm depth (Figure 10A). However,

at the 30-40 cm depth, no environmental factor significantly correlated with bacterial community variation (Figure 10B).

Archaeal community variation at the 10-20 cm depth was significantly correlated to peroxidase ( $p=0.011$ ), sulfatase ( $p=0.026$ ), and  $\beta$ -glucosidase ( $p=0.018$ ) activities, but no other environmental variables had a significant correlation except water table ( $p=0.082$ ), which was marginally significant (Figure 11A). At the 30-40 cm depth, archaeal community variation is not significantly correlated to any environmental factor measured in this study (Figure 11B). The combined variation of axes PCO1 and PCO2 for bacteria at the 10-20 cm depth explained 59.97% of total community variation, while combined axes for bacteria at the 30-40 cm depth explained 76.83% of total community variation. Likewise, combined axes for archaea at the 10-20 cm depth explained 46.09%, while the 30-40 cm depth explained 61.77% of total community variation.

## 1.5 Discussion

### 1.5.1 Plant Functional Group Effects on Bacterial and Archaeal Community Composition

Acidic, oligotrophic bogs were originally considered sterile (Waksman and Stevens, 1929) but have since been found to contain a diverse assemblage of microorganisms. Peat monoliths in our mesocosm experiment were comprised of *Sphagnum*-derived peats with low pH and contained bacterial and archaeal assemblages observed in acid-tolerant bogs (Dedysh et al., 2006) rather than fen systems dominated by neutral pH phylotypes (Lin et al., 2012). Bacterial community composition was dominated by Acidobacteria species regardless of plant functional group treatments or soil depth with the most abundant OTU classified taxonomically as most closely related to *Terriglobus roseus* (7% relative abundance) of the family Acidobacteriaceae. Recent molecular studies have identified that Acidobacteria are best suited to acidic, low-nutrient conditions, being long-lived with the ability to tolerate fluctuations in soil hydration, and function in plant polymer hydrolysis utilizing carbon sources including hemicellulose, cellulose, and chitin (Ward et al., 2009).

Plant functional groups appear to influence certain phyla only within the more shallow 10-20 cm depth, indicating plant traits may have more direct influence on bacterial and archaeal community composition closer to the peat surface. Noticeable differences observed between sedge and ericaceous shrub treatments within the 10-20 cm depth suggest that Ericaceae may suppress the relative abundance of Acidobacteria and promote increased abundance of Bacteroidetes and Gammaproteobacteria taxa. Bacteroidetes species are hypothesized to function mainly in plant polymer hydrolysis (Tveit et al., 2012) and are known to respond in a predictable manner to changes in carbon availability (Fierer et al., 2007). In contrast, sedge treatments were found to promote increased homogeneity with strong

dominance by Acidobacteria. With time, plant functional groups may begin to influence microbial communities at deeper peat depths, but as of two months following vegetation manipulations there has yet to be any significant influences attributed to plant functional group composition.

Certain species of Alpha-, and Gammaproteobacteria as well as Verrucomicrobia are linked to methane oxidizing capabilities under aerobic soil conditions (Dedysh, 2009), which may explain the trends showing increased abundance within the more shallow 10-20 cm depth under sedge treatments whose aerenchyma are supplying the anaerobic peat with increase oxygen availability. Due to high water table conditions at the time of sampling, our chosen sampling depths may be experiencing oxygen limitation within Ericaceae treatments, subsequently limiting the relative abundance of aerobic methane oxidizing bacterial communities initially hypothesized. In the absence of aerenchyma and ebullition, methane created under anaerobic soil conditions must naturally diffuse up through the aerobic layers near the peat surface (0-10 cm depth at time of study) prior to atmospheric release, becoming susceptible to oxidation and subsequent conversion to carbon dioxide (Schlesinger, 1997) by methane oxidizing bacteria. However, current water table conditions at the 10-20 cm depth appear to limit these naturally occurring methane oxidation processes. Subsequently, an increase in available methane under sedge-mediated aerobic rhizosphere conditions near the peat surface may encourage the growth of methane oxidizing bacteria and promote oxidizing redox conditions.

Anaerobic fermentation processes are prevalent functional pathways for carbon mineralization (Tveit et al., 2012), whose waste products via microbial metabolism become the substrates for methanogenic archaea to carry out both hydrogenotrophic and acetoclastic methanogenesis (Brauer et al., 2006; Whalen, 2005; Horn et al., 2003). A lack of distinction in the relative abundance of microbial species between plant functional group treatments within the 30-40 cm depth may be the result of

abrupt changes in aboveground plant biomass, as the system has not regained equilibrium with the environmental conditions. Further work is needed to fully gauge the sustained effects plant functional groups may have on influencing bacterial community structure and function throughout the peat profile.

The dominant methanogenic Euryarchaeota included *Methanomicrobiales* and *Methanobacteriales* especially in the 10-20 cm depth. Both of these archaeal orders are considered hydrogenotrophic, producing CH<sub>4</sub> from H<sub>2</sub> and CO<sub>2</sub>, and their high abundance correlates well with previous studies claiming hydrogenotrophic methanogenesis is the dominant pathway in oligotrophic bogs and other northern peatlands (Horn et al., 2003; Keller and Bridgham, 2007). A large proportion of unclassified Euryarchaeota were observed at the 30-40 cm depth and may be associated with any number of different archaeal orders found in acidic bogs, making predictions about plant functional group influences on archaeal community composition difficult to discern. In addition, these unclassified Euryarchaeota may represent novel lineages not yet described in peatland soils and should be further investigated.

A previous peatland microbial sequencing study (*see* Lin et al., 2012) found members of *Methanosarcinales* were the dominant methanogenic archaea in their bog soils. *Methanosarcinales* are metabolically more diverse than *Methanomicrobiales* and *Methanobacteriales* and can carry out acetoclastic, hydrogenotrophic, and methylotrophic methanogenesis (Garcia et al., 2000). Our results indicated that *Methanosarcinales* were present in relatively low abundances throughout the peat profile, suggesting that acetoclastic methanogenesis was not a dominant methanogenic pathway within these peat soils. The relatively low abundance of *Methanosarcinales* in our study may be explained by the increased abundance of sulfate reducing bacteria (e.g. Deltaproteobacteria taxa) and their seasonal influences on acetate concentrations in the peat soil. Shannon and White (1996) found that

acetate concentrations in bog porewater began to decline in late summer and early fall as sulfate reducers consumed the accumulated acetate. Further analysis of the competition between sulfate reducers and acetoclastic archaea for porewater acetate in early September 2011 could explain the relatively low abundances of *Methanosarcinales* observed in this study.

### *1.5.2 Limitations of DNA-based Sequencing Studies*

Initial changes in the relative recovery of both bacterial and archaeal community composition were observed only in the 10-20 cm depth. These changes may be the direct result of plant functional group manipulations and their subsequent effects on peat soil conditions, but may also be explained in part by the techniques used in this study. High throughput sequencing of extracted soil microbial DNA represents total community composition as it cannot distinguish between viable and non-viable cells (van Elsas and Boersma, 2011) nor dormant versus active cells (Lennon and Jones, 2011). Sequencing data from this study might be representative of microbial community composition prior to plant functional group manipulations as dormant microorganisms can maintain low metabolic activity until environmental conditions become favorable (Jones and Lennon, 2010) and DNA from non-viable microbes can persist in soil environments for considerable periods of time. As such, this study may lack the resolution necessary to accurately quantify the immediate changes in microbial structure following plant community changes. Future studies should strive to sequence both DNA and RNA from environmental samples to capture total community composition and active community composition, respectively. An RNA-based sequencing analysis may provide greater insight into the immediate changes in microbial community structure following plant functional group manipulations as only the active microbial communities at the time of sampling would be detected.



### *1.5.3 Conclusions*

This study revealed the initial response of bacterial and archaeal community composition within an oligotrophic bog to changes in plant functional group dominance and demonstrated distinct microbial responses between sedge and Ericaceae treatments as well as changes in peat depth. Our results suggest that the more shallow 10-20 cm depth is experiencing more pronounced changes in both bacterial and archaeal community structure with the dominant vascular plant functional groups, sedges and Ericaceae, driving opposing effects on microbial community abundance. Sedge-dominated treatments appear to promote aerobic methane oxidizing bacterial communities as a result of increased soil oxygen availability supplied via aerenchyma, while also promoting decreased anaerobic microbial heterogeneity overall. In contrast, Ericaceae treatments under high water table conditions may experience an increase in the relative abundance of anaerobic microbial communities, which suggests that under high water tables Ericaceae may have less direct influence on regulating atmospheric methane flux and possibly encourage decreased plant-based carbon mineralization and function as atmospheric carbon sinks through decreased decomposition.

## **1.6 Acknowledgements**

I thank Carley Kratz from Michigan Technological University for initial guidance with molecular techniques and extracellular enzyme assays. I thank Lynette Potvin and Joe DesRochers from the USDA Forest Service Northern Research Station for technical assistance. I thank Evan Kane and Aleta Daniels from Michigan Technological University for their contributions to porewater chemistry methods and data collection. I also thank Julien Tremblay from the DOE Joint Genome Institute for methods describing the JGI Illumina analysis pipeline. All sequencing services were provided by the DOE Joint Genome Institute, Walnut Creek, California (<http://www.jgi.doe.gov/>). Our research was funded by the National Science Foundation grant DEB-1146149 and by the USDA Forest Service, Northern Research Station.

## **1.7 Literature Cited**

1. Allison, S.D. 2005. Cheaters, diffusion and nutrients constrain decomposition by microbial enzymes in spatially structured environments. *Ecology Letters* 8, 626–635.
2. Andersen, R., Chapman, S.J., Artz, R.R.E. 2012. Microbial communities in natural and disturbed peatlands: A review. *Soil Biology and Biochemistry* 57, 979-994.
3. Artz, R.R.E. 2009. Microbial community structure and carbon substrate use in northern peatlands. *In*: Baird, A.J., Belyea, L.R., Comas, X., Reeve, A.S., Slater, L.D., Eds. *Carbon Cycling in Northern Peatlands*. American Geophysical Union, 111–129.

4. Artz, R.R.E., Anderson, I.C., Chapman, S.J., Hagn, A., Schlöter, M., Potts, J.M., Campbell, C.D. 2007. Changes in fungal community composition in response to vegetational succession during the natural regeneration of cutover peatlands. *Microbial Ecology* 54, 508–522.
5. Bonfante, P., Genre, A. 2010. Mechanisms underlying beneficial plant–fungus interactions in mycorrhizal symbiosis. *Nature Communications* 1, 1–11.
6. Borga, P., Nilsson, M., Tunlid, A. 1994. Bacterial communities in peat in relation to botanical composition as revealed by phospholipid fatty acid analysis. *Soil Biology and Biochemistry* 26, 841–848.
7. Brauer, S.L., Yashiro, E., Ueno, N.G., Yavitt, J.B., Zinder, S.H. 2006. Characterization of acid-tolerant H<sub>2</sub>/CO<sub>2</sub>-utilizing methanogenic enrichment cultures from an acidic peat bog in New York State. *FEMS Microbiology Ecology* 57, 206–216.
8. Breeuwer, A., Robroek, B.J.M., Limpens, J., Heijmans, M.M.P.D., Schouten, M.G.C., Berendse, F. 2009. Decreased summer water table depth affects peatland vegetation. *Basic and Applied Ecology* 10, 330–339.
9. Bridgman, S.D., Pastor, J., Janssens, J.A., Chapin, C., Malterer, T.J. 1996. Multiple limiting gradients in peatlands: a call for a new paradigm. *Wetlands* 16, 45–65.
10. Cadillo-Quiroz, H., Yavitt, J.B., Zinder, S.H., Thies, J.E. 2009. Diversity and community structure of archaea inhabiting the rhizosphere of two contrasting plants from an acidic bog. *Microbial Ecology* 59, 757–767.

11. Cairney, J.W.G., Burke, R.M. 1998. Extracellular enzyme activities of the ericoid mycorrhizal endophyte *Hymenoscyphus ericae* (Read) Korf & Kernan: their likely roles in decomposition of dead plant tissue in soil. *Plant and Soil* 205, 181–192.
12. Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K., Fierer, N., Peña, A.G., Goodrich, J.K., Gordon, J.I., Huttley, G.A., Kelley, S.T., Knights, D., Koenig, J.E., Ley, R.E., Lozupone, C.A., McDonald, D., Muegge, B.D., Pirrung, M., Reeder, J., Sevinsky, J.R., Turnbaugh, P.J., Walters, W.A., Widmann, J., Yatsunenko, T., Zaneveld, J., Knight, R. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7, 335–336.
13. Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer, N., Knight, R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences of the United States of America* 108, 4516–4522.
14. Clymo, R.S. 1984. The limits to peat bog growth. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences* 303, 605–654.
15. Clymo, R.S., Turunen, J., Tolonen, K. 1998. Carbon accumulation in peatland. *Oikos* 368–388.
16. Crum, H. 1991. *A Focus on Peatlands and Peat Mosses*. 4<sup>th</sup> Ed. The University of Michigan Press, Ann Arbor.
17. Dedys, S.N. 2009. Exploring methanotroph diversity in acidic northern wetlands: molecular and cultivation-based studies. *Microbiology* 78, 655–669.

18. Dedys, S.N., Panikov, N.S., Tiedje, J.M. 1998. Acidophilic methanotrophic communities from *Sphagnum* peat bogs. *Applied and Environmental Microbiology* 64, 922–929.
19. De Haan, H., De Boer, T. 1987. Applicability of light absorbance and fluorescence as measures of concentration and molecular size of dissolved organic carbon in humic Lake Tjeukemeer. *Water Research* 21, 731–734.
20. Dinsmore, K.J., Billett, M.F., Skiba, U.M., Rees, R.M., Drewer, J., Helfter, C. 2010. Role of the aquatic pathway in the carbon and greenhouse gas budgets of a peatland catchment. *Global Change Biology* 16, 2750–2762.
21. Farrick, K.K., Price, J.S. 2009. Ericaceous shrubs on abandoned block-cut peatlands: implications for soil water availability and *Sphagnum* restoration. *Ecohydrology* 2, 530–540.
22. Fierer, N., Bradford, M.A., Jackson, R.B. 2007. Toward an ecological classification of soil bacteria. *Ecology* 88, 1354–1364.
23. Fisk, M., Ruether, K.F., Yavitt, J.B. 2003. Microbial activity and functional composition among northern peatland ecosystems. *Soil Biology and Biochemistry* 35, 591–602.
24. Garcia, J.L., Patel, B.K.C., Ollivier, B. 2000. Taxonomic phylogenetic and ecological diversity of methanogenic archaea. *Anaerobe* 6, 205–226.
25. Gorham, E. 1991. Northern peatlands: role in the carbon cycle and probable responses to climatic warming. *Ecological Applications* 1, 182–195.

26. Groisman, P.Y., Knight, R.W., Easterling, D.R., Karl, T.R., Hegerl, G.C., Razuvaev, V.N. 2005. Trends in intense precipitation in the climate record. *Journal of climate* 18, 1326–1350.
27. Hilbert, D.W., Roulet, N.T., Moore, T.R. 2000. Modelling and analysis of peatlands as dynamical systems. *Journal of Ecology* 88, 230–242.
28. Holzapfel-Pschorn, A., Conrad, R., Seiler, W. 1986. Effects of vegetation on the emission of methane from submerged paddy soil. *Plant and Soil* 92, 223–233.
29. Horn, M.A., Matthies, C., Küsel, K., Schramm, A., Drake, H.L. 2003. Hydrogenotrophic methanogenesis by moderately acid-tolerant methanogens of a methane-emitting acidic peat. *Applied and Environmental Microbiology* 69, 74–83.
30. Inderjit, Mallik, A.U. 2002. Can *Kalmia angustifolia* interference to black spruce (*Picea mariana*) be explained by allelopathy? *Forest Ecology and Management* 160, 75–84.
31. Jaatinen, K., Fritze, H., Laine, J., Laiho, R. 2007. Effects of short- and long-term water-level drawdown on the populations and activity of aerobic decomposers in a boreal peatland. *Global Change Biology* 13, 491–510.
32. Joannis, G.D., Bradley, R.L., Preston, C.M., Munson, A.D. 2007. Soil enzyme inhibition by condensed litter tannins may drive ecosystem structure and processes: the case of *Kalmia angustifolia*. *New Phytologist* 175, 535–546.
33. Jones, S.E., Lennon, J.T. 2010. Dormancy contributes to the maintenance of

- microbial diversity. *Proceedings of the National Academy of Sciences* 107, 5881-5886.
34. Keller, J.K., Bridgham, S.D. 2007. Pathways of anaerobic carbon cycling across an ombrotrophic-minerotrophic peatland gradient. *Limnology and Oceanography* 96–107.
35. Kim, S.Y., Lee, S.H., Freeman, C., Fenner, N., Kang, H. 2008. Comparative analysis of soil microbial communities and their responses to the short-term drought in bog, fen, and riparian wetlands. *Soil Biology and Biochemistry* 40, 2874–2880.
36. Kolka, R.K., Sebestyen, S.D., Verry, E.S., Brooks, K.N. 2011. Peatland biogeochemistry and watershed hydrology at the Marcell Experimental Forest. Taylor & Francis Group, LLC, Boca Raton.
37. Laanbroek, H.J. 2009. Methane emission from natural wetlands: interplay between emergent macrophytes and soil microbial processes. A mini-review. *Annals of Botany* 105, 141-153.
38. Lennon, J.T., Jones, S.E. 2011. Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nature Reviews Microbiology* 9, 119-130.
39. Lin, X., Green, S., Tfaily, M.M., Prakash, O., Konstantinidis, K.T., Corbett, J.E., Chanton, J.P., Cooper, W.T., Kostka, J.E. 2012. Microbial community structure and activity linked to contrasting biogeochemical gradients in bog and fen environments of the Glacial Lake Agassiz Peatland. *Applied and Environmental Microbiology* 78, 7023–7031.

40. Lindahl, B.O., Taylor, A.F., Finlay, R.D. 2002. Defining nutritional constraints on carbon cycling in boreal forests – towards a less “phytcentric” perspective. *Plant and Soil* 242, 123–135.
41. Magoc, T., Salzberg, S.L. 2011. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27, 2957–2963.
42. Myers, B., Webster, K.L., McLaughlin, J.W., Basiliko, N. 2011. Microbial activity across a boreal peatland nutrient gradient: the role of fungi and bacteria. *Wetlands Ecology and Management* 20, 77–88.
43. Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Wagner, H. 2011. *vegan: Community Ecology Package*. R package version 1.17-9.
44. Olefeldt, D., Roulet, N.T. 2012. Effects of permafrost and hydrology on the composition and transport of dissolved organic carbon in a subarctic peatland complex. *Journal of Geophysical Research* 117, 1-15.
45. Peltoniemi, K., Fritze, H., Laiho, R. 2009. Response of fungal and actinobacterial communities to water-level drawdown in boreal peatland sites. *Soil Biology and Biochemistry* 41, 1902–1914.
46. Roulet, N.T., Moore, T.R., Bubier, J., Lafleur, P. 1992. Northern fens: methane flux and climatic change. *Tellus B* 44, 100–105.
47. Saiya-Cork, K.R., Sinsabaugh, R.L., Zak, D.R. 2002. The effects of long term nitrogen deposition on extracellular enzyme activity in an *Acer saccharum* forest



- soil. *Soil Biology and Biochemistry* 34, 1309-1315.
48. Schlesinger, W.H. 1997. *Biogeochemistry: An Analysis of Global Change*. 2<sup>nd</sup> Edition. Academic Press, San Diego, California, USA.
49. Shannon, R.D., White, J.R. 1996. The effects of spatial and temporal variations in acetate and sulfate on methane cycling in two Michigan peatlands. *Limnology and Oceanography* 41, 435-443.
50. Sowerby, A., Emmett, B., Beier, C., Tietema, A., Peñuelas, J., Estiarte, M., Van Meeteren, M.J.M., Hughes, S., Freeman, C. 2005. Microbial community changes in heathland soil communities along a geographical gradient: interaction with climate change manipulations. *Soil Biology and Biochemistry* 37, 1805–1813.
51. Strack, M., Waller, M.F., Waddington, J.M. 2006. Sedge succession and peatland methane dynamics: a potential feedback to climate change. *Ecosystems* 9, 278–287.
52. Straková, P., Niemi, R.M., Freeman, C., Peltoniemi, K., Toberman, H., Heiskanen, I., Fritze, H., Laiho, R. 2011. Litter type affects the activity of aerobic decomposers in a boreal peatland more than site nutrient and water table regimes. *Biogeosciences* 8, 2741–2755.
53. Sundh, I., Nilsson, M., Granberg, G., Svensson, B.H. 1994. Depth distribution of microbial production and oxidation of methane in northern boreal peatlands. *Microbial Ecology* 27, 253–265.
54. Trettin, C.C., Laiho, R., Minkkinen, K., Laine, J. 2006. Influence of climate change factors on carbon dynamics in northern forested peatlands. *Canadian*

- Journal of Soil Science 86, 269–280.
55. Tsonis, A.A., Triantafyllou, G.N., Georgakakos, K.P. 1996. Hydrological applications of satellite data: 1. Rainfall estimation. *Journal of Geophysical Research* 101, 26517–26525.
56. Tveit, A., Schwacke, R., Svenning, M.M., Urich, T. 2012. Organic carbon transformations in high-Arctic peat soils: key functions and microorganisms. *The ISME Journal* (2012) 1-13.
57. van Elsas, J.D., Boersma, F.G.H. 2011. A review of molecular methods to study the microbiota of soil and the mycosphere. *European Journal of Soil Biology* 47, 77–87.
58. Waksman, S.A., Stevens, K.R. 1929. Contribution to the chemical composition of peat. V. The role of microorganisms in peat formation and decomposition. *Soil Science* 28, 315-339.
59. Ward, N.L., Challacombe, J.F., Janssen, P.H., Henrissat, B., Coutinho, P.M., Wu, M., Xie, G., Haft, D.H., Sait, M., Badger, J., Barabote, R.D., Bradley, B., Brettin, T.S., Brinkac, L.M., Bruce, D., Creasy, T., Daugherty, S.C., Davidsen, T.M., DeBoy, R.T., Detter, J.C., Dodson, R.J., Durkin, A.S., Ganapathy, A., Gwinn-Giglio, M., Han, C.S., Khouiri, H., Kiss, H., Kothari, S.P., Madupu, R., Nelson, K.E., Nelson, W.C., Paulsen, I., Penn, K., Ren, Q., Rosovitz, M.J., Selengut, J.D., Shrivastava, S., Sullivan, S.A., Tapia, R., Thompson, L.S., Watkins, K.L., Yang, Q., Yu, C., Zafar, N., Zhou, L., Kuske, C.R. 2009. Three genomes from the phylum *Acidobacteria* provide insight into the lifestyle of these microorganisms in soils. *Applied and Environmental Microbiology* 75, 2046-2056.

60. Whalen, S.C. 2005. Biogeochemistry of methane exchange between natural wetlands and the atmosphere. *Environmental Engineering Science* 22, 73-94.
61. Weishaar, J.L., Aiken, G.R., Bergamaschi, B.A., Fram, M.S., Fujii, R., Mopper, K. 2003. Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon. *Environmental Science & Technology* 37, 4702-4708.
62. Weltzin, J.F., Pastor, J., Harth, C., Bridgham, S.D., Updegraff, K., Chapin, C.T. 2000. Response of bog and fen plant communities to warming and water-table manipulations. *Ecology* 81, 3464–3478.

## 1.8 Tables

**Table 1** Oxidative and hydrolytic extracellular enzymes assayed in bulk peat at the 10-20 cm depth with corresponding enzyme commission number (EC), role in nutrient cycling, and substrate.

Enzyme	Abbreviation	EC	Type	Nutrient Cycle	Substrate
Phenol oxidase	POX	1.10.3.2	Oxidative	Carbon	L-DOPA
Peroxidase	PER	1.11.1.7	Oxidative	Carbon	L-DOPA
Sulfatase	SULF	3.1.6.1	Hydrolytic	Sulfur	4-MUB-sulfate potassium salt
$\beta$ -1,4-glucosidase	GLUC	3.2.1.21	Hydrolytic	Carbon	4-MUB- $\beta$ -D-glucopyranoside
Acid-phosphatase	PHOS	3.1.3.2	Hydrolytic	Phosphorus	4-MUB-phosphate

L-DOPA = L-3,4-dihydroxyphenylalanine; 4-MUB = 4-methylumbelliferyl

**Table 2** Bacterial and archaeal diversity indicated by Chao1 richness and equitability evenness in plant functional group treatments.

Treatment	Depth (cm)	Bacteria			Archaea		
		Sequence			Sequence		
		Read	Chao1	Equitability	Read	Chao1	Equitability
Sedge	10-20	197,708	2,082 ± 363	0.78 ± 0.03	965	10 ± 5	0.79 ± 0.08
Sedge	30-40	291,657	1,968 ± 174	0.74 ± 0.02	2,471	11 ± 3	0.80 ± 0.15
Unmanipulated	10-20	276,899	2,123 ± 197	0.77 ± 0.03	976	8 ± 4	0.74 ± 0.11
Unmanipulated	30-40	224,881	1,948 ± 218	0.75 ± 0.04	2,309	12 ± 3	0.80 ± 0.12
Ericaceae	10-20	253,350	2,038 ± 265	0.75 ± 0.03	804	13 ± 11	0.73 ± 0.17
Ericaceae	30-40	290,087	1,920 ± 262	0.75 ± 0.04	2,188	11 ± 5	0.84 ± 0.16

Calculations based on OTU tables rarefied to the smallest total sequencing read per sample to minimize the effects of variable sampling effort in comparing diversity across samples. Sequence reads are the sum of sequences generated per treatment combination. Diversity values are arranged by treatment combination with numbers following "±" indicating standard deviation.

**Table 3** Mean ( $\pm$  SE) percent relative abundance of bacterial phyla and archaeal orders between plant functional groups within sampling depths.

Bacteria	10-20 cm Depth			30-40 cm Depth		
	Ericaceae	Unmanipulated	Sedge	Ericaceae	Unmanipulated	Sedge
Phylum	(%)	(%)	(%)	(%)	(%)	(%)
Acidobacteria	26 $\pm$ 3	32 $\pm$ 3	36 $\pm$ 1	45 $\pm$ 2	44 $\pm$ 1	48 $\pm$ 1
Bacteroidetes	23 $\pm$ 3	18 $\pm$ 3	13 $\pm$ 2	6 $\pm$ 2	4 $\pm$ 1	2 $\pm$ 0
Bacteria Unknown	14 $\pm$ 1	16 $\pm$ 1	17 $\pm$ 1	19 $\pm$ 0	19 $\pm$ 1	21 $\pm$ 1
Alphaproteobacteria	9 $\pm$ 0	10 $\pm$ 0	11 $\pm$ 1	8 $\pm$ 1	9 $\pm$ 1	8 $\pm$ 0
Gammaproteobacteria	9 $\pm$ 1	6 $\pm$ 0	7 $\pm$ 1	4 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0
Actinobacteria	4 $\pm$ 0	4 $\pm$ 0	3 $\pm$ 0	3 $\pm$ 0	4 $\pm$ 1	3 $\pm$ 0
Verrucomicrobia	3 $\pm$ 0	3 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0	4 $\pm$ 0
Cyanobacteria	5 $\pm$ 2	3 $\pm$ 1	2 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 0
Deltaproteobacteria	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0	3 $\pm$ 0	3 $\pm$ 0
Planctomycetes	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0	3 $\pm$ 0	2 $\pm$ 0
Firmicutes	1 $\pm$ 0	1 $\pm$ 0	1 $\pm$ 0	3 $\pm$ 1	3 $\pm$ 1	3 $\pm$ 0
Other	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0	2 $\pm$ 0

Archaea	10-20 cm Depth			30-40 cm Depth		
	Ericaceae	Unmanipulated	Sedge	Ericaceae	Unmanipulated	Sedge
Order	(%)	(%)	(%)	(%)	(%)	(%)
<i>Cenarchaeales</i>	33 $\pm$ 12	37 $\pm$ 9	21 $\pm$ 8	28 $\pm$ 5	27 $\pm$ 6	24 $\pm$ 5
<i>Methanobacteriales</i>	31 $\pm$ 12	33 $\pm$ 8	37 $\pm$ 7	18 $\pm$ 10	14 $\pm$ 5	6 $\pm$ 3
<i>Methanomicrobiales</i>	22 $\pm$ 9	18 $\pm$ 9	19 $\pm$ 7	1 $\pm$ 1	5 $\pm$ 2	3 $\pm$ 2
Euryarchaeota Unclassified	7 $\pm$ 2	8 $\pm$ 3	15 $\pm$ 5	23 $\pm$ 6	35 $\pm$ 10	43 $\pm$ 9
<i>Halobacteriales</i>	6 $\pm$ 5	2 $\pm$ 1	1 $\pm$ 1	2 $\pm$ 1	0	2 $\pm$ 1
Archaea Unknown	0	2 $\pm$ 1	1 $\pm$ 1	10 $\pm$ 3	8 $\pm$ 2	12 $\pm$ 2
<i>Thermococcales</i>	1 $\pm$ 1	1 $\pm$ 1	1 $\pm$ 1	7 $\pm$ 2	5 $\pm$ 1	3 $\pm$ 2
Crenarchaeota Unclassified	1 $\pm$ 1	0	0	0	0	0
<i>Thaumarchaeota</i>	0	0	2 $\pm$ 1	7 $\pm$ 2	4 $\pm$ 2	3 $\pm$ 1
<i>Thermoproteales</i>	0	0	0	1 $\pm$ 1	1 $\pm$ 1	1 $\pm$ 1
<i>Methanocellales</i>	0	0	0	0	0	1 $\pm$ 1
<i>Methanosarcinales</i>	0	0	2 $\pm$ 2	3 $\pm$ 2	2 $\pm$ 2	2 $\pm$ 2

Numbers following " $\pm$ " indicate standard error. "Other" contains the seven relatively rare remaining phyla and three Proteobacteria classes that combined represent less than 5% of total community composition.

**Table 4** Peat soil biogeochemistry, relative percentage of vegetation, and extracellular enzyme activities by plant functional group treatments within depths.

Treatment	Depth (cm)	pH	DOC (mg L <sup>-1</sup> )	TDN (mg L <sup>-1</sup> )	E <sub>2</sub> :E <sub>3</sub>	SUVA <sub>254</sub>
Sedge	10-20	3.96 ± 0.18	92.6 ± 19.8	2.9 ± 1.0	7.0 ± 0.7	4.5 ± 0.2
Sedge	30-40	3.84 ± 0.11	99.7 ± 23.0	2.9 ± 1.0	6.4 ± 0.7	4.3 ± 0.2
Unmanipulated	10-20	3.82 ± 0.20	111.6 ± 21.4	2.9 ± 0.6	6.5 ± 0.8	4.2 ± 0.2
Unmanipulated	30-40	3.83 ± 0.13	117.5 ± 22.7	3.1 ± 0.7	5.9 ± 0.7	4.0 ± 0.2
Ericaceae	10-20	3.86 ± 0.11	111.3 ± 25.0	2.9 ± 0.7	6.8 ± 0.8	4.0 ± 0.2
Ericaceae	30-40	3.77 ± 0.08	121.5 ± 23.5	3.3 ± 0.8	5.9 ± 0.7	3.8 ± 0.2

Treatment	Water Table (cm)	Relative Percentage (%)				
		<i>Carex oligosperma</i>	<i>Andromeda glaucophylla</i>	<i>Chamaedaphne calyculata</i>	<i>Kalmia polifolia</i>	<i>Vaccinium oxycoccos</i>
Sedge	-7.4 ± 3.2	29 ± 1	0 ± 1	1 ± 8	0 ± 11	6 ± 11
Unmanipulated	-7.4 ± 2.4	14 ± 15	0 ± 0	20 ± 1	18 ± 1	34 ± 5
Ericaceae	-7.2 ± 2.5	0 ± 10	0 ± 0	23 ± 9	21 ± 4	42 ± 8

Treatment	Depth (cm)	Extracellular Enzyme Activity				
		Phenol Oxidase (μmol h <sup>-1</sup> g <sup>-1</sup> )	Peroxidase (μmol h <sup>-1</sup> g <sup>-1</sup> )	β-Glucosidase (nmol hr <sup>-1</sup> g <sup>-1</sup> )	Sulfatase (nmol hr <sup>-1</sup> g <sup>-1</sup> )	Acid-Phosphatase (nmol hr <sup>-1</sup> g <sup>-1</sup> )
Sedge	10-20	1.42 ± 1.93	6.90 ± 4.47	2630.6 ± 866.2	35.4 ± 36.1	6588.8 ± 1425.6
Ericaceae	10-20	0.92 ± 2.21	8.45 ± 5.65	3059.8 ± 948.9	11.2 ± 9.0	5194.6 ± 2237.1
Unmanipulated	10-20	0.62 ± 1.01	6.80 ± 5.16	2970.7 ± 1037.6	13.5 ± 13.3	6241.6 ± 2383.9

Numbers following "±" indicate standard deviation


**Table 5** Statistical results for envfit function utilizing environmental data for bacteria and archaea at both 10-20 cm and 30-40 cm depths.

	Bacteria				Archaea			
	10-20 cm		30-40 cm		10-20 cm		30-40 cm	
	r <sup>2</sup>	p-value	r <sup>2</sup>	p-value	r <sup>2</sup>	p-value	r <sup>2</sup>	p-value
pH	0.02	0.858	0.02	0.843	0.15	0.204	0.02	0.829
Water Table	0.07	0.492	0.02	0.790	0.24	0.082	0.01	0.883
<i>Carex</i> %	0.52	<b>0.001*</b>	0.19	0.117	0.02	0.872	0.13	0.254
<i>Chamae</i> %	0.14	0.233	0.07	0.486	0.01	0.920	0.06	0.567
<i>Kalmia</i> %	0.12	0.301	0.14	0.218	0.03	0.782	0.06	0.546
<i>Vaccin</i> %	0.25	0.085	0.16	0.169	0.00	0.965	0.16	0.199
DOC	0.26	<b>0.040*</b>	0.07	0.505	0.11	0.333	0.09	0.388
TDN	0.07	0.491	0.06	0.543	0.10	0.401	0.10	0.318
E <sub>2</sub> :E <sub>3</sub>	0.27	<b>0.034*</b>	0.16	0.178	0.15	0.214	0.16	0.166
SUVA <sub>254</sub>	0.21	0.099	0.05	0.621	0.17	0.147	0.08	0.392
Phenol Oxidase	0.04	0.735	-	-	0.06	0.561	-	-
Peroxidase	0.20	0.135	-	-	0.41	<b>0.011*</b>	-	-
Sulfatase	0.61	<b>0.002*</b>	-	-	0.32	<b>0.026*</b>	-	-
β-Glucosidase	0.26	0.058	-	-	0.35	<b>0.018*</b>	-	-
Acid-Phosphatase	0.03	0.782	-	-	0.00	0.969	-	-


\*p-value < 0.05




## 1.9 Figures




**Figure 1** Rank abundance curves for relative abundance of A) bacterial and B) archaeal OTUs obtained from Illumina sequencing. Graph axes are displayed in log-log scale for clarity.




**Figure 2** Water table depth in mesocosm bins for high water table treatment (HWT) and low water table treatment (LWT) during the two-week peat coring period.




**Figure 3** Bacteria phylum-level community composition between sampling depth. “Other” contains seven relatively rare remaining phyla and three Proteobacteria classes that represent less than 5% of total community composition.




**Figure 4** Bacteria phylum-level community composition between plant functional group treatments within sampling depths. “Other” contains seven relatively rare remaining phyla and three Proteobacteria classes that represent less than 5% of total community composition




**Figure 5** Archaea order-level community composition between sampling depth.



**Figure 6** Archaea order-level community composition between plant functional group treatments within sampling depths.




**Figure 7** Heatmap visualization for top 100 most abundant bacterial OTUs indicating relative abundance by depth. Refer to Supplemental Table S1 for taxonomic identification of relevant OTUs.




**Figure 8** Heatmap visualization for all archaeal OTUs indicating relative abundance by depth. Refer to Supplemental Table S3 for taxonomic identification of relevant OTUs.









**Figure 10** PCoA of top 100 OTUs with envfit environmental vectors for A) bacteria at 10-20 cm depth and B) bacteria at 30-40 cm depth. Environmental vectors that had a significant ( $p < 0.05$ ) influence on microbial community composition within the ordination are plotted in red.



**Figure 11** PCoA of all OTUs with envfit environmental vectors for A) archaea at 10-20 cm depth and B) archaea at 30-40 cm depth. Environmental vectors that had a significant ( $p < 0.05$ ) influence on microbial community composition within the ordination are plotted in red.



**Figure 12** Mean relative abundance of all methanotrophic bacteria summed between sampling depths. Error bars represent standard error.



**Figure 13** Mean relative abundance of all methanogenic archaea summed between sampling depths. Error bars represent standard error.

## 1.10 Supplemental

**Table S1** Depth effects on top 50 most abundant bacterial OTUs as calculated through the OTU Category Significance ANOVA in QIIME.


OTU	Total Read Sequences	Proportion within study (%)	Raw p-value	Benferroni (corrected)	FDR	10-20cm (mean)	30-40cm (mean)	Consensus Lineage
0*	109,909	7.16	0.000	0.000	0.000	0.0332	0.1937	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae; g_Terriglobus; s_Terriglobus roseus;
1*	52,291	3.41	0.000	0.000	0.000	0.0357	0.0820	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae;
3*	46,882	3.06	0.000	0.008	0.000	0.0983	0.0069	k_Bacteria; p_Bacteroidetes;
2*	42,675	2.78	0.000	0.000	0.000	0.1034	0.0140	k_Bacteria; p_Bacteroidetes; c_Sphingobacteriales; o_Sphingobacteriales;
6	39,416	2.57	0.304	15.187	0.330	0.0472	0.0522	k_Bacteria; p_Proteobacteria; c_Alphaproteobacteria; o_Rhodospirillales; f_Rhodospirillaceae; g_Rhodospirillum; s_Rhodospirillum elegans;
5*	31,541	2.06	0.000	0.000	0.000	0.0139	0.0570	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae; g_Terriglobus; s_Terriglobus roseus;
9*	28,740	1.87	0.000	0.008	0.000	0.0291	0.0411	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae; g_Terriglobus; s_Terriglobus roseus;
17*	25,932	1.69	0.000	0.001	0.000	0.0514	0.0202	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae; g_Terriglobus; s_Terriglobus roseus;
20*	24,235	1.58	0.000	0.006	0.000	0.0438	0.0228	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae; g_Terriglobus; s_Terriglobus roseus;
13	23,858	1.55	0.002	0.123	0.004	0.0520	0.0201	k_Bacteria; p_Bacteroidetes; c_Sphingobacteriales; o_Sphingobacteriales;
10*	23,092	1.50	0.000	0.000	0.000	0.0590	0.0063	k_Bacteria; p_Proteobacteria; c_Gammaproteobacteria;
11*	21,683	1.41	0.000	0.001	0.000	0.0085	0.0358	k_Bacteria;
7	20,104	1.31	0.241	12.034	0.274	0.0330	0.0244	k_Bacteria; p_Proteobacteria; c_Gammaproteobacteria;
16	20,010	1.30	0.158	7.894	0.184	0.0224	0.0264	k_Bacteria; p_Acidobacteria; c_Acidobacteria; o_Solihacteriales; f_Solihacteriaceae; g_Candidatus Solihacter; s_Candidatus Solihacter usitatus;
24*	18,879	1.23	0.000	0.000	0.000	0.0127	0.0295	k_Bacteria; p_Firmicutes; c_Clostridia;
4*	18,223	1.19	0.000	0.008	0.000	0.0030	0.0323	k_Bacteria; p_Acidobacteria; c_Acidobacteria; o_Solihacteriales; f_Solihacteriaceae; g_Candidatus Solihacter; s_Candidatus Solihacter usitatus;
21*	17,534	1.14	0.000	0.000	0.000	0.0113	0.0290	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae; g_Terriglobus; s_Terriglobus roseus;
58	14,625	0.95	0.007	0.336	0.009	0.0244	0.0161	k_Bacteria; p_Cyanobacteria; c_Oscillatoriales; f_Oscillatoriaceae; g_Merococcus; s_Merococcus chthonoplastes;
8	12,163	0.79	0.005	0.255	0.007	0.0239	0.0064	k_Bacteria; p_Bacteroidetes; c_Sphingobacteriales; o_Sphingobacteriales;
18	12,096	0.79	0.250	12.513	0.278	0.0210	0.0065	k_Bacteria; p_Proteobacteria; c_Alphaproteobacteria; o_Rhodospirillales; f_Rhodospirillaceae; g_Terriglobus; s_Terriglobus roseus;
22	12,025	0.78	0.020	0.976	0.025	0.0109	0.0187	k_Bacteria; p_Acidobacteria; c_Acidobacteria (class); o_Acidobacteriales; f_Acidobacteriaceae; g_Terriglobus; s_Terriglobus roseus;
34	11,610	0.76	0.021	1.039	0.025	0.0172	0.0134	k_Bacteria; p_Cyanobacteria;
15*	10,714	0.70	0.000	0.015	0.001	0.0238	0.0098	k_Bacteria; p_Acidobacteria; c_Acidobacteria; o_Solihacteriales; f_Solihacteriaceae; g_Candidatus Solihacter; s_Candidatus Solihacter usitatus;
23*	9,913	0.64	0.000	0.000	0.000	0.0047	0.0164	k_Bacteria;
14*	9,485	0.62	0.000	0.000	0.000	0.0047	0.0164	k_Bacteria; p_Acidobacteria; c_Acidobacteria; o_Solihacteriales; f_Solihacteriaceae; g_Candidatus Solihacter; s_Candidatus Solihacter usitatus;
57	9,273	0.60	0.357	17.835	0.379	0.0138	0.0112	k_Bacteria; p_Firmicutes; c_Clostridia;
20*	9,127	0.59	0.000	0.000	0.000	0.0038	0.0154	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
72*	9,039	0.59	0.000	0.000	0.000	0.0016	0.0070	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
47*	8,937	0.59	0.000	0.003	0.000	0.0016	0.0083	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
43*	8,804	0.55	0.000	0.018	0.001	0.0075	0.0024	k_Bacteria; p_Proteobacteria; c_Gammaproteobacteria;
32*	8,415	0.55	0.000	0.020	0.001	0.0153	0.0085	k_Bacteria; p_Acidobacteria; c_Acidobacteria; o_Solihacteriales; f_Solihacteriaceae; g_Candidatus Solihacter; s_Candidatus Solihacter usitatus;
108*	7,877	0.51	0.000	0.000	0.000	0.0048	0.0123	k_Bacteria; p_Firmicutes; c_Clostridia;
29*	7,806	0.51	0.000	0.010	0.000	0.0066	0.0145	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
19	7,666	0.50	0.008	0.419	0.011	0.0151	0.0080	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
33	7,355	0.49	0.845	42.258	0.845	0.0089	0.0092	k_Bacteria; p_Acinobacteria; c_Acinobacteria (class); o_Acinobacteriales; f_Acinobacteriaceae; g_Acinobacterium;
46*	6,790	0.44	0.000	0.000	0.000	0.0148	0.0044	k_Bacteria; p_Acinobacteria; c_Acinobacteria (class); o_Acinobacteriales; f_Acinobacteriaceae; g_Acinobacterium;
63*	6,702	0.44	0.000	0.016	0.001	0.0153	0.0042	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
42*	6,652	0.43	0.000	0.000	0.000	0.0040	0.0115	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
26*	6,625	0.43	0.000	0.000	0.000	0.0040	0.0115	k_Bacteria; p_Verrucomicrobia; c_Verrucomicrobiales; f_Verrucomicrobiaceae; g_Verrucomicrobia; s_Verrucomicrobia;
27*	6,222	0.41	0.000	0.001	0.000	0.0147	0.0038	k_Bacteria; p_Proteobacteria; c_Deltaproteobacteria; o_Deltaproteobacterales; f_Geobacteriaceae; g_Geobacter; s_Geobacter propionicus;
79*	6,180	0.40	0.000	0.000	0.000	0.0165	0.0022	k_Bacteria;
31	6,152	0.40	0.579	28.965	0.591	0.0074	0.0080	k_Bacteria; p_Proteobacteria; c_Alphaproteobacteria; o_Rhodospirillales; f_Rhodospirillaceae; g_Acinobacteriaceae;
36	6,115	0.40	0.427	21.349	0.445	0.0084	0.0071	k_Bacteria; p_Acinobacteria; c_Acinobacteria; o_Solihacteriales; f_Solihacteriaceae; g_Candidatus Solihacter; s_Candidatus Solihacter usitatus;
30	5,719	0.37	0.008	0.399	0.011	0.0055	0.0089	k_Bacteria;
25	5,409	0.35	0.020	1.005	0.025	0.0036	0.0076	k_Bacteria;
4246*	5,115	0.33	0.000	0.000	0.000	0.0022	0.0087	k_Bacteria;
52*	5,095	0.33	0.000	0.000	0.000	0.0108	0.0021	k_Bacteria; p_Proteobacteria; c_Alphaproteobacteria; o_Rhodospirillales; f_Rhodospirillaceae; g_Acinobacteriaceae;
55	5,070	0.33	0.003	0.139	0.004	0.0035	0.0093	k_Bacteria; p_Planctomycetes; c_Planctomycetes; o_Gemmatimonadetes; f_Isoptomycetaceae; g_Singulisphaera; s_Singulisphaera acidiphila;
92	5,013	0.33	0.020	0.989	0.025	0.0041	0.0070	k_Bacteria; p_Acinobacteria; c_Acinobacteria; o_Solihacteriales; f_Solihacteriaceae; g_Candidatus Solihacter; s_Candidatus Solihacter usitatus;
40	4,677	0.30	0.003	0.145	0.004	0.0037	0.0067	k_Bacteria; p_Acinobacteria; c_Acinobacteria (class);

\*p<0.05 following Benferroni Correction

**Table S2** Depth effects on archaeal OTUs as calculated through the OTU Category Significance ANOVA in QIIME and list of all archaeal OTUs with consensus lineage from the oligotrophic bog mesocosm study.


OTU	Total Read Sequences	Proportion within study (%)	Consensus Lineage
<b>45*</b>	2,513	25.87	k__Archaea;p__Euryarchaeota;c__Thaumarchaeota;o__Methanobacteriales;f__Methanobacterium;Methanobacterium beijingense;
<b>103*</b>	1,475	15.19	k__Archaea;p__Euryarchaeota;c__Thaumarchaeota;o__Methanobacteriales;f__Methanobacterium;Methanobacterium beijingense;
<b>139*</b>	1,281	13.19	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;o__Cenarchaeales;f__Cenarchaeaceae;g__Nitrosopumilus;Nitrosopumilus maritimus;
234	1,001	10.31	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;o__Cenarchaeales;f__Cenarchaeaceae;
<b>303*</b>	806	8.30	k__Archaea;
<b>239*</b>	520	5.35	k__Archaea;p__Euryarchaeota;c__Methanomicrobiales;f__Methanomicrobiales;g__Methanofolius;Methanofolius limitans;
<b>312*</b>	405	4.17	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;
225	372	3.83	k__Archaea;p__Euryarchaeota;
235	320	3.29	k__Archaea;p__Euryarchaeota;c__Thermococci;o__Thermococcales;f__Thermococcales;g__Thermococcus;
391	226	2.33	k__Archaea;p__Euryarchaeota;c__Methanomicrobiales;o__Methanosarcinales;f__Methanosarcinaceae;g__Methanosarcina;
<b>441*</b>	141	1.45	k__Archaea;p__Euryarchaeota;
658	89	0.92	k__Archaea;p__Euryarchaeota;
1143	75	0.77	k__Archaea;p__Crenarchaeota;c__Thermoprotei;o__Thermoproteales;
863	67	0.69	k__Archaea;p__Euryarchaeota;c__Methanobacteriales;f__Methanobacteriales;g__Methanobacterium;
795	51	0.53	k__Archaea;p__Euryarchaeota;
843	37	0.38	k__Archaea;p__Euryarchaeota;c__Methanomicrobiales;o__Methanosarcinales;f__Methanosarcinaceae;g__Methanosarcina thermophila;
1341	33	0.34	k__Archaea;p__Euryarchaeota;c__Halobacteriales;f__Halobacteriales;g__Halobacterium;
1868	24	0.25	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;
4501	22	0.23	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;
1419	21	0.22	k__Archaea;p__Euryarchaeota;
1587	19	0.20	k__Archaea;p__Euryarchaeota;c__Thermococci;o__Thermococcales;f__Thermococcales;g__Thermococcus;
2594	18	0.19	k__Archaea;p__Euryarchaeota;c__Halobacteriales;f__Halobacteriales;g__Halobacteriaceae;
3290	18	0.19	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;
3917	18	0.19	k__Archaea;p__Euryarchaeota;c__Methanobacteriales;o__Methanobacteriales;f__Methanobacterium;Methanobacterium beijingense;
1281	18	0.19	k__Archaea;p__Euryarchaeota;c__Methanomicrobiales;o__Methanocellales;f__Methanocellaceae;g__Methanocella paludicola;
2595	17	0.18	k__Archaea;p__Euryarchaeota;c__Methanomicrobiales;f__Methanomicrobiales;g__Methanofolius;Methanofolius limitans;
3145	13	0.13	k__Archaea;p__Euryarchaeota;c__Halobacteriales;f__Halobacteriales;
1418	13	0.13	k__Archaea;p__Euryarchaeota;c__Thermococci;o__Thermococcales;f__Thermococcales;g__Thermococcus;
2593	9	0.09	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;o__Cenarchaeales;f__Cenarchaeaceae;
3140	9	0.09	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;o__Cenarchaeales;f__Cenarchaeaceae;g__Nitrosopumilus;Nitrosopumilus maritimus;
3141	9	0.09	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;o__Nitrososphaerales;f__Nitrososphaeraeae;g__Candidatus Nitrososphaera;
4507	9	0.09	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;
4509	9	0.09	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;g__Halobacterium;
1869	8	0.08	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;
3144	8	0.08	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;g__Halobacterium;Halobacterium alkalicum;
4506	8	0.08	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;g__Halobacterium;Halobacterium lipolyticum;
2258	6	0.06	k__Archaea;p__Crenarchaeota;
4503	6	0.06	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;g__Halobacterium;
3143	5	0.05	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;g__Haladaptatus;Haladaptatus paucihalophilus;
4510	4	0.04	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;
4500	3	0.03	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;
4508	3	0.03	k__Archaea;p__Euryarchaeota;c__Halobacteriales;o__Halobacteriales;f__Halobacteriaceae;
3147	2	0.02	k__Archaea;p__Crenarchaeota;
5647	2	0.02	k__Archaea;p__Euryarchaeota;c__Methanomicrobiales;f__Methanomicrobiales;g__Methanomicrobiales;

\*p<0.05 following Bonferroni Correction.




**Figure S1** Electrophoresis DNA quantity visualization on 2:1% agarose gel for DNA extractions at the 10-20 cm depth.






**Figure S2** Electrophoresis DNA quantity visualization on 2:1% agarose gel for DNA extractions at the 30-40 cm depth.



**Figure S3** Relative abundance of all known methanotrophic bacteria within plant functional group treatments between sampling depths as found in this study. AP=Alphaproteobacteria, GP=Gammaproteobacteria, Verr=Verrucomicrobia.



**Figure S4** Mean relative abundance of all known methanogenic archaea plus unclassified Euryarchaeota within plant functional group treatments between sampling depths as found in this study.